

**NOVEL CHROMOPHORES/FLUOROPHORES AND
METHODS FOR USING THE SAME**CROSS REFERENCE TO RELATED APPLICATIONS

5 This application is a continuation-in-part of Application Serial No. PCT/US00/28477
filed on October 13, 2000; which application is a continuation-in-part of the following
applications: application serial no. 09/418,529 filed October 14, 1999; application serial
no. 09/418,917 filed October 15, 1999; application serial no. 09/418,922 filed October 15,
1999; application serial no. 09/444,338 filed November 19, 1999; application serial no.
10 09/444,341 filed November 19, 1999; application serial no. 09/457,556 filed December 9,
1999; application serial no. 09/458,477 filed December 9, 1999; application serial no.
09/458,144 filed December 9, 1999; and application serial no. 09/457,898 filed December
9, 1999; all of which applications claim priority to application serial no. 09/210,330 filed
December 11, 1998; as well as application serial no. 60/211,627 filed on June 14, 2000;
5 application serial no. 60/211,687 filed on June 14, 2000; application serial no. 60/211,609
filed on June 14, 2000; application serial no. 60/211,626 filed on June 14, 2000;
application serial no. 60/211,880 filed on June 14, 2000; application serial no. 60/211,607
filed on June 14, 2000; application serial no. 60/211,766 filed on June 14, 2000;
application serial no. 60/211,888 filed on June 14, 2000; and application serial no.
20 60/212,070 filed on June 14, 2000; as well as International Application Serial No.
PCT/US99/29405 filed December 10, 1999, which application claims priority to application
serial no. 09/210,330 filed December 11, 1998; the disclosures of which application are
incorporated in their entirety herein.

INTRODUCTIONField of the Invention

The field of this invention is chromoproteins and fluorescent proteins.

Background of the Invention

30 Labeling is a tool for marking a protein, cell, or organism of interest and plays a
prominent role in many biochemistry, molecular biology and medical diagnostic
applications. A variety of different labels have been developed, including radiolabels,
chromolabels, fluorescent labels, chemiluminescent labels, etc. However, there is
continued interest in the development of new labels. Of particular interest is the
development of new protein labels, including chromo- and/or fluorescent protein labels.

Relevant Literature

35 U.S. Patents of interest include: 6,066,476; 6,020,192; 5,985,577; 5,976,796;
5,968,750; 5,968,738; 5,958,713; 5,919,445; 5,874,304; and 5,491,084. International
Patent Publications of interest include: WO 00/46233; WO 99/49019; and DE 197 18 640
A. Also of interest are: Anderluh et al., Biochemical and Biophysical Research

Communications (1996) 220:437-442; Dove et al., Biological Bulletin (1995) 189:288-297; Fradkov et al., FEBS Lett. (2000) 479(3):127-30; Gurskaya et al., FEBS Lett., (2001) 507(1):16-20; Gurskaya et al., BMC Biochem. (2001) 2:6; Lukyanov, K., et al (2000) J Biol Chemistry 275(34):25879-25882; Macek et al., Eur. J. Biochem. (1995) 234:329-335; Martynov et al., J Biol Chem. (2001) 276:21012-6; Matz, M.V., *et al.* (1999) Nature Biotechnol., 17:969-973; Tersikh et al., Science (2000) 290:1585-8; Tsien, Annual Rev. of Biochemistry (1998) 67:509-544; Tsien, Nat. Biotech. (1999) 17:956-957; Ward et al., J. Biol. Chem. (1979) 254:781-788; Wiedermann et al., Jahrestagung der Deutschen Gesellschaft fur Tropenokologie-gto. Ulm. 17-19.02.1999. Poster P-4.20; Yarbrough et al., Proc Natl Acad Sci U S A (2001) 98:462-7.

SUMMARY OF THE INVENTION

Nucleic acid compositions encoding novel chromo/fluoroproteins and mutants thereof, as well as the proteins encoded the same, are provided. The proteins of interest are proteins that are colored and/or fluorescent, where this feature arises from the interaction of two or more residues of the protein. The subject proteins are further characterized in that they are either obtained from non-bioluminescent Cnidarian, e.g., Anthozoan, species or are obtained from Anthozoan non-Pennatulacean (sea pen) species. Specific proteins of interest include proteins obtained from the following specific Anthozoan species: Anemonia majano (NFP-1), Clavularia sp. (NFP-2), Zoanthus sp. (NFP-3 & NFP-4), Discosoma striata (NFP-5), Discosoma sp. "red"(NFP-6), Anemonia sulcata (NFP-7), Discosoma sp "green" (NFP-8), and Discosoma sp."magenta" (NFP-9). Also of interest are proteins that are substantially similar to, or mutants of, the above specific proteins. Also provided are fragments of the nucleic acids and the peptides encoded thereby, as well as antibodies to the subject proteins and transgenic cells and organisms. The subject protein and nucleic acid compositions find use in a variety of different applications. Finally, kits for use in such applications, e.g., that include the subject nucleic acid compositions, are provided.

BREIF DESCRIPTION OF THE FIGURES

Figure 1 provides the nucleotide and amino acid sequence of wild type amFP486 (NFP-1). (SEQ ID NO:01 & 02)

Figure 2 provides the nucleotide and amino acid sequence of wild type cFP484 (NFP-2). (SEQ ID NO:03 & 04)

Figure 3 provides the nucleotide and amino acid sequence of wild type zFP506 (NFP-3). (SEQ ID NO:05 & 06)

Figure 4 provides the nucleotide and amino acid sequence of wild type zFP538 (NFP-4). (SEQ ID NO:07 & 08)

Figure 5 provides the nucleotide and amino acid sequence of wild type dsFP483 (NFP-5). (SEQ ID NO:09 & 10)

Figure 6 provides the nucleotide and amino acid sequence of wild type drFP583 (NFP-6). (SEQ ID NO:11 & 12); as well as the nucleotide and amino acid sequence of an alternative version thereof.

Figure 7 provides the nucleotide and amino acid sequence of wild type asFP600 (NFP-7). (SEQ ID NO:13 & 14)

Figure 8 provides the nucleotide and amino acid sequence of wild type dgFP512 (NFP-8). (SEQ ID NO:15 & 16)

Figure 9 provides the nucleotide and amino acid sequence of wild type dmFP592 (NFP-9). (SEQ ID NO:17 & 18)

Figure 10 provides the nucleotide and amino acid sequence of mut32-NA (SEQ ID NO: 27 & 28)

Figure 11 provides the nucleotide and amino acid sequence of FP3-NA (SEQ ID NO:29 & 30)

Figure 12 provides the amino acid sequence of FP3-yellow (SEQ ID NO:31)

Figure 13 provides the amino acid sequence of FP3-Yellow bright (SEQ ID NO:32)

Figure 14 provides the nucleotide and amino acid sequence of NFP4-NA (SEQ ID NO:33 & 34).

Figure 15 provides additional sequence information of NFP-6 mutants.

Figure 16 provides the nucleotide sequence of humanized NFP-6. (SEQ ID NO:35)

Figure 17 provides the nucleotide sequence of mutant E5-NA. (SEQ ID NO:36)

Figure 18 provides the nucleotide sequence of mutant E57. (SEQ ID NO:37)

Figure 19 provides the nucleotide sequence of mutant E57-NA (SEQ ID NO:38)

Figure 20 provides the nucleotide and amino acid sequence of mutant FP7-NA (SEQ ID NO:39 & 40).

Figure 21 provides the nucleotide and amino acid sequence of humanized FP7. (SEQ ID NO:41 & 42).

Figure 22 provides the nucleotide and amino acid sequence of humanized 6/9Q (SEQ ID NO:43 & 44).

DEFINITIONS

In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Maniatis, Fritsch & Sambrook, "Molecular Cloning: A Laboratory Manual (1982); "DNA Cloning: A Practical Approach," Volumes I and II (D.N. Glover ed. 1985); "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid Hybridization" (B.D. Hames & S.J. Higgins eds. (1985)); "Transcription and Translation" (B.D. Hames & S.J. Higgins eds. (1984)); "Animal Cell

Culture" (R.I. Freshney, ed. (1986)); "Immobilized Cells and Enzymes" (IRL Press, (1986)); B. Perbal, "A Practical Guide To Molecular Cloning" (1984).

A "vector" is a replicon, such as plasmid, phage or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment.

A "DNA molecule" refers to the polymeric form of deoxyribonucleotides (adenine, guanine, thymine, or cytosine) in either single stranded form or a double-stranded helix. This term refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, inter alia, in linear DNA molecules (e.g., restriction fragments), viruses, plasmids, and chromosomes.

A DNA "coding sequence" is a DNA sequence which is transcribed and translated into a polypeptide *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences from eukaryotic (e.g., mammalian) DNA, and synthetic DNA sequences. A polyadenylation signal and transcription termination sequence may be located 3' to the coding sequence.

As used herein, the term "hybridization" refers to the process of association of two nucleic acid strands to form an antiparallel duplex stabilized by means of hydrogen bonding between residues of the opposite nucleic acid strands.

The term "oligonucleotide" refers to a short (under 100 bases in length) nucleic acid molecule.

"DNA regulatory sequences", as used herein, are transcriptional and translational control sequences, such as promoters, enhancers, polyadenylation signals, terminators, and the like, that provide for and/or regulate expression of a coding sequence in a host cell.

A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site, as well as protein binding domains responsible for the binding of RNA polymerase. Eukaryotic promoters will often, but not always, contain "TATA" boxes and "CAT" boxes. Various promoters, including inducible promoters, may be used to drive the various vectors of the present invention.

As used herein, the terms "restriction endonucleases" and "restriction enzymes" refer to bacterial enzymes, each of which cut double-stranded DNA at or near a specific nucleotide sequence.

A cell has been "transformed" or "transfected" by exogenous or heterologous DNA when such DNA has been introduced inside the cell. The transforming DNA may or may not be integrated (covalently linked) into the genome of the cell. In prokaryotes, yeast, and mammalian cells for example, the transforming DNA may be maintained on an episomal element such as a plasmid. With respect to eukaryotic cells, a stably transformed cell is one in which the transforming DNA has become integrated into a chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter cells containing the transforming DNA. A "clone" is a population of cells derived from a single cell or common ancestor by mitosis. A "cell line" is a clone of a primary cell that is capable of stable growth *in vitro* for many generations.

A "heterologous" region of the DNA construct is an identifiable segment of DNA within a larger DNA molecule that is not found in association with the larger molecule in nature. Thus, when the heterologous region encodes a mammalian gene, the gene will usually be flanked by DNA that does not flank the mammalian genomic DNA in the genome of the source organism. In another example, heterologous DNA includes coding sequence in a construct where portions of genes from two different sources have been brought together so as to produce a fusion protein product. Allelic variations or naturally-occurring mutational events do not give rise to a heterologous region of DNA as defined herein.

As used herein, the term "reporter gene" refers to a coding sequence attached to heterologous promoter or enhancer elements and whose product may be assayed easily and quantifiably when the construct is introduced into tissues or cells.

The amino acids described herein are preferred to be in the "L" isomeric form. The amino acid sequences are given in one-letter code (A: alanine; C: cysteine; D: aspartic acid; E: glutamic acid; F: phenylalanine; G: glycine; H: histidine; I: isoleucine; K: lysine; L: leucine; M: methionine; N: asparagine; P: proline; Q: glutamine; R: arginine; S: serine; T: threonine; V: valine; W: tryptophan; Y: tyrosine; X: any residue). NH₂ refers to the free amino group present at the amino terminus of a polypeptide. COOH refers to the free carboxy group present at the carboxy terminus of a polypeptide. In keeping with standard polypeptide nomenclature, *J Biol. Chem.*, 243 (1969), 3552-59 is used.

The term "immunologically active" defines the capability of the natural, recombinant or synthetic chromo/fluorescent protein, or any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies. As

used herein, "antigenic amino acid sequence" means an amino acid sequence that, either alone or in association with a carrier molecule, can elicit an antibody response in a mammal. The term "specific binding," in the context of antibody binding to an antigen, is a term well understood in the art and refers to binding of an antibody to the antigen to which the antibody was raised, but not other, unrelated antigens.

As used herein the term "isolated" is meant to describe a polynucleotide, a polypeptide, an antibody, or a host cell that is in an environment different from that in which the polynucleotide, the polypeptide, the antibody, or the host cell naturally occurs.

Bioluminescence (BL) is defined as emission of light by living organisms that is well visible in the dark and affects visual behavior of animals (See e.g., Harvey, E. N. (1952). *Bioluminescence*. New York: Academic Press; Hastings, J. W. (1995). *Bioluminescence*. In: *Cell Physiology* (ed. by N. Speralakis). pp. 651-681. New York: Academic Press.; Wilson, T. and Hastings, J. W. (1998). *Bioluminescence. Annu Rev Cell Dev Biol* **14**, 197-230.). Bioluminescence does not include so-called ultra-weak light emission, which can be detected in virtually all living structures using sensitive luminometric equipment (Murphy, M. E. and Sies, H.(1990). Visible-range low-level chemiluminescence in biological systems. *Meth.Enzymol.* **186**, 595-610; Radotic, K, Radenovic, C, Jeremic, M. (1998.) Spontaneous ultra-weak bioluminescence in plants: origin, mechanisms and properties. *Gen Physiol Biophys* **17**, 289-308), and from weak light emission which most probably does not play any ecological role, such as the glowing of bamboo growth cone (Totsune, H., Nakano, M., Inaba, H.(1993). Chemiluminescence from bamboo shoot cut. *Biochem. Biophys.Res Comm.* **194**, 1025-1029) or emission of light during fertilization of animal eggs (Klebanoff, S. J., Froeder, C. A., Eddy, E. M., Shapiro, B. M. (1979). Metabolic similarities between fertilization and phagocytosis. Conservation of peroxidatic mechanism. *J. Exp. Med.* **149**, 938-953; Schomer, B. and Epel, D. (1998). Redox changes during fertilization and maturation of marine invertebrate eggs. *Dev Biol* **203**, 1-11).

DESCRIPTION OF THE SPECIFIC EMBODIMENTS

Nucleic acid compositions encoding novel chromo/fluoroproteins and mutants thereof, as well as the proteins encoded the same, are provided. The proteins of interest are proteins that are colored and/or fluorescent, where this feature arises from the interaction of two or more residues of the protein. The subject proteins are further characterized in that they are either obtained from non-bioluminescent Cnidarian, e.g., Anthozoan, species or are obtained from non-Pennatulacean (sea pen) Anthozoan species. Specific proteins of interest include proteins obtained from the following specific Anthozoan species: *Anemonia majano* (NFP-1), *Clavularia* sp. (NFP-2), *Zoanthus* sp. (NFP-3 & NFP-4), *Discosoma striata* (NFP-5), *Discosoma* sp. "red"(NFP-6), *Anemonia*

sulcata (NFP-7), *Discosoma* sp “green” (NFP-8), and *Discosoma* sp. “magenta” (NFP-9). Also of interest are proteins that are substantially similar to, or mutants of, the above specific proteins. Also provided are fragments of the nucleic acids and the peptides encoded thereby, as well as antibodies to the subject proteins, and transgenic cells and organisms that include the subject nucleic acid/protein compositions. The subject protein and nucleic acid compositions find use in a variety of different applications. Finally, kits for use in such applications, e.g., that include the subject nucleic acid compositions, are provided.

Before the subject invention is described further, it is to be understood that the invention is not limited to the particular embodiments of the invention described below, as variations of the particular embodiments may be made and still fall within the scope of the appended claims. It is also to be understood that the terminology employed is for the purpose of describing particular embodiments, and is not intended to be limiting. Instead, the scope of the present invention will be established by the appended claims.

In this specification and the appended claims, the singular forms “a,” “an” and “the” include plural reference unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs.

Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range, and any other stated or intervening value in that stated range, is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges, and are also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in the invention.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs. Although any methods, devices and materials similar or equivalent to those described herein can be used in the practice or testing of the invention, the preferred methods, devices and materials are now described.

All publications mentioned herein are incorporated herein by reference for the purpose of describing and disclosing the cell lines, vectors, methodologies and other

invention components that are described in the publications which might be used in connection with the presently described invention.

In further describing the subject invention, the subject nucleic acid compositions will be described first, followed by a discussion of the subject protein compositions, antibody compositions and transgenic cells/organisms. Next a review of representative methods in which the subject proteins find use is provided.

NUCLEIC ACID COMPOSITIONS

As summarized above, the subject invention provides nucleic acid compositions encoding chromo- and fluoroproteins and mutants thereof, as well as fragments and homologues of these proteins. By chromo and/or fluorescent protein is meant a protein that is colored, i.e., is pigmented, where the protein may or may not be fluorescent, e.g., it may exhibit low, medium or high fluorescence upon irradiation with light of an excitation wavelength. In any event, the subject proteins of interest are those in which the colored characteristic, i.e., the chromo and/or fluorescent characteristic, is one that arises from the interaction of two or more residues of the protein, and not from a single residue, more specifically a single side chain of a single residue, of the protein. As such, fluorescent proteins of the subject invention do not include proteins that exhibit fluorescence only from residues that act by themselves as intrinsic fluors, i.e., tryptophan, tyrosine and phenylalanine. As such, the fluorescent proteins of the subject invention are fluorescent proteins whose fluorescence arises from some structure in the protein that is other than the above specified single residues, e.g., it arises from an interaction of two or more residues.

By nucleic acid composition is meant a composition comprising a sequence of DNA having an open reading frame that encodes a chromo/fluoro polypeptide of the subject invention, i.e., a chromo/fluoroprotein gene, and is capable, under appropriate conditions, of being expressed as a chromo/fluoro protein according to the subject invention. Also encompassed in this term are nucleic acids that are homologous, substantially similar or identical to the nucleic acids of the present invention. Thus, the subject invention provides genes and coding sequences thereof encoding the proteins of the subject invention, as well as homologs thereof. The subject nucleic acids are present in other than their natural environment, e.g., they are isolated, present in enriched amounts, etc., from their naturally occurring environment, e.g., the organism from which they are obtained.

The nucleic acids are further characterized in that they encode proteins that are either from: (1) non-bioluminescent species, often non-bioluminescent Cnidarian species, e.g., non-bioluminescent Anthozoan species; or (2) from Anthozoan species that are not Pennatulacean species, i.e., that are not sea pens. As such, the nucleic acids may

encode proteins from bioluminescent Anthozoan species, so long as these species are not Pennatulacean species, e.g., that are not Renillan or Ptilosarcan species. Specific nucleic acid compositions of interest are those that encode proteins (and mutants thereof) from the following specific Anthozoan species: *Anemonia majano*, *Clavularia* sp., *Zoanthus* sp., *Discosoma striata*, *Discosoma* sp. "red", *Anemonia sulcata*, *Discosoma* sp. "green", and *Discosoma* sp. "magenta." Each of these particular types of nucleic acid compositions of interest is now discussed in greater detail individually.

Anemonia majano (NFP-1; AmCyan; RCFP-1)

In these embodiments, the nucleic acid compositions are found in, derived from, or are mutants, homologues of, a nucleic acid found in an organism from the Anthozoan Sub-class Zoantharia, often Order Actiniaria, more often Sub-order Endomyaria, usually Family Actiniidae, and more usually Genus *Anemonia*, where in many embodiments, the organism is *Anemonia majano*, where the specific wild type protein of interest is amFP486 (i.e., NFP-1; RCFP-1). The wild type cDNA coding sequence for amFP486 is provided in SEQ ID NO: 01. In addition to nucleic acids encoding the wild type protein and fragments thereof, also of interest are nucleic acids that encode homologues and mutants of the wild type protein. Specific mutants of interest include, but are not limited to: Mut15, Mut32, and FP1-NA (a non-aggregating mutant), where these specific mutants are further described in the experimental section, *infra*.

Clavularia sp. (NFP-2; RCPF-2)

In these embodiments, the nucleic acid compositions are found in, derived from, or are mutants or homologues of, nucleic acids found in Anthozoan organisms from Sub-class Alcyonaria, often Order Stolonifera, and more often the Family Clavulariidae, where the organism is usually from the Genus *Clavularia*, and in certain embodiments, the organism is *Clavularia* sp., where the specific wild type fluorescent protein of interest is cFP484 (i.e., NFP-2; RCPF-2). The wild type cDNA coding sequence for cFP484 is provided in SEQ ID No: 03. In addition to nucleic acids encoding the wild type sequence and fragments thereof, also of interest are nucleic acids that encode homologues and mutants of the wild type protein. Specific mutants of interest include, but are not limited to: $\Delta 19$ cFP484 and $\Delta 38$ cFP484, where these specific mutants are further described in the experimental section, *infra*.

Zoanthus sp. I (NFP-3; ZsGreen; RCFP-3)

In these embodiments, the nucleic acid compositions are found in, derived from, or are mutants or homologues of, nucleic acids found in Anthozoan organisms from Sub-class Zoantharia, often Order Zoanthidea, more often Sub-order Brachycnemia, usually Family Zoanthidae, and more usually Genus Zoanthus, where in certain embodiments, the organism is *Zoanthus sp.*, where the specific wild type fluorescent protein of interest is zFP506 (i.e., NFP-3; RCFP-3). The wild type cDNA coding sequence for zFP506 is provided in SEQ ID Nos: 05. In addition to nucleic acids encoding the wild type sequence and fragments thereof, also of interest are nucleic acids that encode homologues and mutants of the wild type protein. Specific mutants of interest include, but are not limited to: N66M; NFP-3NA (a non-aggregating mutant); yellow; yellow bright, etc., where these specific mutants are further described in the experimental section, infra.

Zoanthus sp. II (NFP-4; ZsYellow; RCFP-4)

In these embodiments, the nucleic acid compositions are found in, derived from, or are mutants or homologues of, nucleic acids found in Anthozoan organisms from Sub-class Zoantharia, often Order Zoanthidea, more often Sub-order Brachycnemia, usually Family Zoanthidae, and more usually Genus Zoanthus, where in certain embodiments, the organism is *Zoanthus sp.*, where the specific wild type fluorescent protein of interest is zFP538 (NFP-4; RCFP-4). The wild type cDNA coding sequence for zFP538 is provided in SEQ ID NO. 07. In addition to nucleic acids encoding the wild type sequence and fragments thereof, also of interest are nucleic acids that encode homologues and mutants of the wild type protein. Specific mutants of interest include, but are not limited to: M129V; FP4-NA (a non-aggregating mutant); green; etc., where these specific mutants are further described in the experimental section, infra.

Discosoma striata (NFP-5; RCFP-5)

In these embodiments, the nucleic acid compositions are found in, derived from, or are mutants or homologues of, nucleic acids found in Anthozoan organisms from Sub-class Zoantharia, often Order Corallimopharia, more often Family Discosomatidae, and usually Genus Discosoma, where in certain embodiments, the organism is *Discosoma striata*, where the specific wild type fluorescent protein of interest is dsFP483 (NFP-5; RCFP-5). The wild type cDNA coding sequence for dsFP483 is provided in SEQ ID NO: 09. In addition to nucleic acids encoding the wild type sequence and fragments thereof, also of interest are nucleic acids that encode homologues and mutants of the wild type protein.

Discosoma sp. "red" (NFP-6; RCFP-6; DsRed; DsRed2)

In these embodiments, the nucleic acid compositions are found in, derived from, or are mutants or homologues of, nucleic acids found in Anthozoan organisms from Sub-class Zoantharia, often Order Corallimopharia, more often Family Discosomatidae, and usually Genus *Discosoma*, where in certain embodiments, the organism is *Discosoma sp. "red"*, where the specific wild type fluorescent protein of interest is drFP583 (NFP-6; RCFP-6). The wild type cDNA coding sequence for drFP583 is provided in SEQ ID NO: 11. In addition to nucleic acids encoding the wild type sequence and fragments thereof, also of interest are nucleic acids that encode homologues and mutants of the wild type protein. Specific mutants of interest include, but are not limited to: E5, E5-NA (a non-aggregating mutant); E8, E5up, E5down, E57, FP6-NA (a non-aggregating mutant), AG4, AG45, E83, 6/9 Q, 6/9 Q-NA, 6/92G; etc., where these specific mutants are further described in the experimental section, infra.

Anemonia sulcata (NFP-7; AsRed; RCFP-7)

In these embodiments, the nucleic acid compositions are found in, derived from, or are mutants or homologues of, nucleic acids found in Anthozoan organisms from Sub-class Zoantharia, often Order Actiniaria, more often Sub-Order Endomyaria, usually Family Actiniidae, and more usually Genus *Anemonia*, e.g., where in certain embodiments the organism is *Anemonia sulcata*, where the specific wild type fluorescent protein of interest is asFP600 (NFP-7; RCFP-7). The wild type cDNA coding sequence for asFP600 is provided in SEQ ID NO:14. In addition to nucleic acids encoding the wild type sequence and fragments thereof, also of interest are nucleic acids that encode homologues and mutants of the wild type protein. Specific mutants of interest include, but are not limited to: Mut1; Mut35-5/Mut1; FP7-NA (a non-aggregating mutant), etc., where these specific mutants are further described in the experimental section, infra.

Discosoma sp "green" (NFP-8; RCFP-8)

In these embodiments, the nucleic acid compositions are found in, derived from, or are mutants or homologues of, nucleic acids found in Anthozoan organisms from Sub-class Zoantharia, often Order Corallimopharia, more often Family Discosomatidae, and usually Genus *Discosoma*, where in certain embodiments, the organism is *Discosoma sp. "green"*, where the specific wild type fluorescent protein of interest is dgFP512 (NFP-8; RCFP-8). The wild type cDNA coding sequence for dgFP512 is provided in SEQ ID NO: 15. In addition to nucleic acids encoding the wild type sequence and fragments thereof,

also of interest are nucleic acids that encode homologues and mutants of the wild type protein.

5 *Discosoma* sp. "magenta" (NFP-9; RCFP-9)

10 In these embodiments, the nucleic acid compositions are found in, derived from, or are mutants or homologues of, nucleic acids found in Anthozoan organisms from Sub-class Zoantharia, often Order Corallimopharia, more often Family Discosomatidae, and usually Genus *Discosoma* where in certain embodiments, the organism is *Discosoma* sp. "magenta", where the specific wild type fluorescent protein of interest is dmFP592 (NFP-9; RCFP-9). The wild type cDNA coding sequence for dmFP592 is provided in SEQ ID NO: 17. In addition to nucleic acids encoding the wild type sequence and fragments thereof, also of interest are nucleic acids that encode homologues and mutants of the wild type protein.

20 In addition to the above described specific nucleic acid compositions, also of interest are homologues of the above sequences. With respect to homologues of the subject nucleic acids, the source of homologous genes may be any species of plant or animal or the sequence may be wholly or partially synthetic. In certain embodiments, sequence similarity between homologues is at least about 20%, sometimes at least about 25 %, and may be 30 %, 35%, 40%, 50%, 60%, 70% or higher, including 75%, 80%, 85%, 90% and 95% or higher. Sequence similarity is calculated based on a reference
25 sequence, which may be a subset of a larger sequence, such as a conserved motif, coding region, flanking region, etc. A reference sequence will usually be at least about 18 nt long, more usually at least about 30 nt long, and may extend to the complete sequence that is being compared. Algorithms for sequence analysis are known in the art, such as BLAST, described in Altschul *et al.* (1990), *J. Mol. Biol.* 215:403-10 (using default settings,
30 i.e. parameters $w=4$ and $T=17$). The sequences provided herein are essential for recognizing related and homologous nucleic acids in database searches. Of particular interest in certain embodiments are nucleic acids of substantially the same length as the nucleic acid identified as SEQ ID NOS: 01, 03, 05, 07, 09, 11, 13, 15, or 17, where by substantially the same length is meant that any difference in length does not exceed
35 about 20 number %, usually does not exceed about 10 number % and more usually does not exceed about 5 number %; and have sequence identity to any of these sequences of at least about 90%, usually at least about 95% and more usually at least about 99% over the entire length of the nucleic acid. In many embodiments, the nucleic acids have a sequence that is substantially similar (i.e. the same as) or identical to the sequences of

SEQ ID NOS: 01, 03, 05, 07, 09, 11, 13, 15, or 17. By substantially similar is meant that sequence identity will generally be at least about 60%, usually at least about 75% and often at least about 80, 85, 90, or even 95%.

Also provided are nucleic acids that encode the proteins encoded by the above described nucleic acids, but differ in sequence from the above described nucleic acids due to the degeneracy of the genetic code.

Also provided are nucleic acids that hybridize to the above described nucleic acid under stringent conditions. An example of stringent hybridization conditions is hybridization at 50°C or higher and 0.1×SSC (15 mM sodium chloride/1.5 mM sodium citrate). Another example of stringent hybridization conditions is overnight incubation at 42°C in a solution: 50 % formamide, 5 × SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5 × Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1 × SSC at about 65°C. Stringent hybridization conditions are hybridization conditions that are at least as stringent as the above representative conditions, where conditions are considered to be at least as stringent if they are at least about 80% as stringent, typically at least about 90% as stringent as the above specific stringent conditions. Other stringent hybridization conditions are known in the art and may also be employed to identify nucleic acids of this particular embodiment of the invention.

Nucleic acids encoding mutants of the proteins of the invention are also provided. Mutant nucleic acids can be generated by random mutagenesis or targeted mutagenesis, using well-known techniques which are routine in the art. In some embodiments, chromo- or fluorescent proteins encoded by nucleic acids encoding homologues or mutants have the same fluorescent properties as the wild-type fluorescent protein. In other embodiments, homologue or mutant nucleic acids encode chromo- or fluorescent proteins with altered spectral properties, as described in more detail herein.

One category of mutant that is of particular interest is the non-aggregating mutant. In many embodiments, the non-aggregating mutant differs from the wild type sequence by a mutation in the N-terminus that modulates the charges appearing on side groups of the N-terminus residues, e.g., to reverse or neutralize the charge, in a manner sufficient to produce a non-aggregating mutant of the naturally occurring protein or mutant, where a particular protein is considered to be non-aggregating if it is determined be non-aggregating using the assay reported in U.S. Patent Application serial no. 60/270,983, the disclosure of which is herein incorporated by reference. More specifically, basic residues located near the N-termini of the proteins are substituted, e.g., Lys and Arg residues close to the N-terminus are substituted with negatively charged or neutral residues. Specific non-aggregating mutants of interest include, but are not limited to: FP1-NA; FP3-NA; FP4-NA; FP6-NA; E5-NA; 6/9Q-NA; 7A-NA; and the like, where these particular non-aggregating mutants are further described infra.

Another category of mutant of particular interest is the modulated oligomerization mutant. A mutant is considered to be a modulated oligomerization mutant if its oligomerization properties are different as compared to the wild type protein. For example, if a particular mutant oligomerizes to a greater or lesser extent than the wild type, it is considered to be an oligomerization mutant. Of particular interest are oligomerization mutants that do not oligomerize, i.e., are monomers under physiological (e.g., intracellular) conditions, or oligomerize to a lesser extent than the wild type, e.g., are dimers or trimers under intracellular conditions.

Nucleic acids of the subject invention may be cDNA or genomic DNA or a fragment thereof. In certain embodiments, the nucleic acids of the subject invention include one or more of the open reading frames encoding specific fluorescent proteins and polypeptides, and introns, as well as adjacent 5' and 3' non-coding nucleotide sequences involved in the regulation of expression, up to about 20 kb beyond the coding region, but possibly further in either direction. The subject nucleic acids may be introduced into an appropriate vector for extrachromosomal maintenance or for integration into a host genome, as described in greater detail below.

The term "cDNA" as used herein is intended to include all nucleic acids that share the arrangement of sequence elements found in native mature mRNA species, where sequence elements are exons and 5' and 3' non-coding regions. Normally mRNA species have contiguous exons, with the intervening introns, when present, being removed by nuclear RNA splicing, to create a continuous open reading frame encoding the protein.

A genomic sequence of interest comprises the nucleic acid present between the initiation codon and the stop codon, as defined in the listed sequences, including all of the introns that are normally present in a native chromosome. It may further include 5' and 3' un-translated regions found in the mature mRNA. It may further include specific transcriptional and translational regulatory sequences, such as promoters, enhancers, etc., including about 1 kb, but possibly more, of flanking genomic DNA at either the 5' or 3' end of the transcribed region. The genomic DNA may be isolated as a fragment of 100 kbp or smaller; and substantially free of flanking chromosomal sequence. The genomic DNA flanking the coding region, either 3' or 5', or internal regulatory sequences as sometimes found in introns, contains sequences required for proper tissue and stage specific expression.

The nucleic acid compositions of the subject invention may encode all or a part of the subject proteins. Double or single stranded fragments may be obtained from the DNA sequence by chemically synthesizing oligonucleotides in accordance with conventional methods, by restriction enzyme digestion, by PCR amplification, etc. For the most part, DNA fragments will be of at least about 15 nt, usually at least about 18 nt or about 25 nt, and may be at least about 50 nt. In some embodiments, the subject nucleic acid

molecules may be about 100 nt, about 200 nt, about 300 nt, about 400 nt, about 500 nt, about 600 nt, about 700 nt, or about 720 nt in length. The subject nucleic acids may encode fragments of the subject proteins or the full-length proteins, e.g., the subject nucleic acids may encode polypeptides of about 25 aa, about 50 aa, about 75 aa, about 100 aa, about 125 aa, about 150 aa, about 200 aa, about 210 aa, about 220 aa, about 230 aa, or about 240 aa, up to the entire protein.

The subject nucleic acids are isolated and obtained in substantial purity, generally as other than an intact chromosome. Usually, the DNA will be obtained substantially free of other nucleic acid sequences that do not include a nucleic acid of the subject invention or fragment thereof, generally being at least about 50%, usually at least about 90% pure and are typically "recombinant", *i.e.* flanked by one or more nucleotides with which it is not normally associated on a naturally occurring chromosome.

The subject polynucleotides (e.g., a polynucleotide having a sequence of SEQ ID NOS: 01 to 17 etc.), the corresponding cDNA, the full-length gene and constructs of the subject polynucleotides are provided. These molecules can be generated synthetically by a number of different protocols known to those of skill in the art. Appropriate polynucleotide constructs are purified using standard recombinant DNA techniques as described in, for example, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., (1989) Cold Spring Harbor Press, Cold Spring Harbor, NY, and under current regulations described in United States Dept. of HHS, National Institute of Health (NIH) Guidelines for Recombinant DNA Research.

Also provided are nucleic acids that encode fusion proteins of the subject proteins, or fragments thereof, which are fused to a second protein, e.g., a degradation sequence, a signal peptide, etc. Fusion proteins may comprise a subject polypeptide, or fragment thereof, and a non-Anthozoan polypeptide (“the fusion partner”) fused in-frame at the N-terminus and/or C-terminus of the subject polypeptide. Fusion partners include, but are not limited to, polypeptides that can bind antibody specific to the fusion partner (e.g., epitope tags); antibodies or binding fragments thereof; polypeptides that provide a catalytic function or induce a cellular response; ligands or receptors or mimetics thereof; and the like. In such fusion proteins, the fusion partner is generally not naturally associated with the subject Anthozoan portion of the fusion protein, and is typically not an Anthozoan protein or derivative/fragment thereof, i.e., it is not found in Anthozoan species.

Also provided are constructs comprising the subject nucleic acids inserted into a vector, where such constructs may be used for a number of different applications, including propagation, protein production, etc. Viral and non-viral vectors may be prepared and used, including plasmids. The choice of vector will depend on the type of cell in which propagation is desired and the purpose of propagation. Certain vectors are useful for amplifying and making large amounts of the desired DNA sequence. Other

domains, usually at least about 8 amino acids in length, more usually at least about 15 amino acids in length, to about 25 amino acids, and up to the complete open reading frame of the gene. After introduction of the DNA, the cells containing the construct may be selected by means of a selectable marker, the cells expanded and then used for expression.

The above described expression systems may be employed with prokaryotes or eukaryotes in accordance with conventional ways, depending upon the purpose for expression. For large scale production of the protein, a unicellular organism, such as *E. coli*, *B. subtilis*, *S. cerevisiae*, insect cells in combination with baculovirus vectors, or cells of a higher organism such as vertebrates, e.g. COS 7 cells, HEK 293, CHO, Xenopus Oocytes, etc., may be used as the expression host cells. In some situations, it is desirable to express the gene in eukaryotic cells, where the expressed protein will benefit from native folding and post-translational modifications. Small peptides can also be synthesized in the laboratory. Polypeptides that are subsets of the complete protein sequence may be used to identify and investigate parts of the protein important for function.

Specific expression systems of interest include bacterial, yeast, insect cell and mammalian cell derived expression systems. Representative systems from each of these categories is are provided below:

Bacteria. Expression systems in bacteria include those described in Chang *et al.*, *Nature* (1978) 275:615; Goeddel *et al.*, *Nature* (1979) 281:544; Goeddel *et al.*, *Nucleic Acids Res.* (1980) 8:4057; EP 0 036,776; U.S. Patent No. 4,551,433; DeBoer *et al.*, *Proc. Natl. Acad. Sci. (USA)* (1983) 80:21-25; and Siebenlist *et al.*, *Cell* (1980) 20:269.

Yeast. Expression systems in yeast include those described in Hinnen *et al.*, *Proc. Natl. Acad. Sci. (USA)* (1978) 75:1929; Ito *et al.*, *J. Bacteriol.* (1983) 153:163; Kurtz *et al.*, *Mol. Cell. Biol.* (1986) 6:142; Kunze *et al.*, *J. Basic Microbiol.* (1985) 25:141; Gleeson *et al.*, *J. Gen. Microbiol.* (1986) 132:3459; Roggenkamp *et al.*, *Mol. Gen. Genet.* (1986) 202:302; Das *et al.*, *J. Bacteriol.* (1984) 158:1165; De Louvencourt *et al.*, *J. Bacteriol.* (1983) 154:737; Van den Berg *et al.*, *Bio/Technology* (1990) 8:135; Kunze *et al.*, *J. Basic Microbiol.* (1985) 25:141; Cregg *et al.*, *Mol. Cell. Biol.* (1985) 5:3376; U.S. Patent Nos. 4,837,148 and 4,929,555; Beach and Nurse, *Nature* (1981) 300:706; Davidow *et al.*, *Curr. Genet.* (1985) 10:380; Gaillardin *et al.*, *Curr. Genet.* (1985) 10:49; Ballance *et al.*, *Biochem. Biophys. Res. Commun.* (1983) 112:284-289; Tilburn *et al.*, *Gene* (1983) 26:205-221; Yelton *et al.*, *Proc. Natl. Acad. Sci. (USA)* (1984) 81:1470-1474; Kelly and Hynes, *EMBO J.* (1985) 4:475479; EP 0 244,234; and WO 91/00357.

Insect Cells. Expression of heterologous genes in insects is accomplished as described in U.S. Patent No. 4,745,051; Friesen *et al.*, "The Regulation of Baculovirus Gene Expression", in: *The Molecular Biology Of Baculoviruses* (1986) (W. Doerfler, ed.); EP 0 127,839; EP 0 155,476; and Vlak *et al.*, *J. Gen. Virol.* (1988) 69:765-776; Miller *et*

al., *Ann. Rev. Microbiol.* (1988) 42:177; Carbonell et al., *Gene* (1988) 73:409; Maeda et al., *Nature* (1985) 315:592-594; Lebacqz-Verheyden et al., *Mol. Cell. Biol.* (1988) 8:3129; Smith et al., *Proc. Natl. Acad. Sci. (USA)* (1985) 82:8844; Miyajima et al., *Gene* (1987) 58:273; and Martin et al., *DNA* (1988) 7:99. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts are described in Luckow et al., *Bio/Technology* (1988) 6:47-55, Miller et al., *Generic Engineering* (1986) 8:277-279, and Maeda et al., *Nature* (1985) 315:592-594.

Mammalian Cells. Mammalian expression is accomplished as described in Dijkema et al., *EMBO J.* (1985) 4:761, Gorman et al., *Proc. Natl. Acad. Sci. (USA)* (1982) 79:6777, Boshart et al., *Cell* (1985) 41:521 and U.S. Patent No. 4,399,216. Other features of mammalian expression are facilitated as described in Ham and Wallace, *Meth. Enz.* (1979) 58:44, Barnes and Sato, *Anal. Biochem.* (1980) 102:255, U.S. Patent Nos. 4,767,704, 4,657,866, 4,927,762, 4,560,655, WO 90/103430, WO 87/00195, and U.S. RE 30,985.

When any of the above host cells, or other appropriate host cells or organisms, are used to replicate and/or express the polynucleotides or nucleic acids of the invention, the resulting replicated nucleic acid, RNA, expressed protein or polypeptide, is within the scope of the invention as a product of the host cell or organism. The product is recovered by any appropriate means known in the art.

Once the gene corresponding to a selected polynucleotide is identified, its expression can be regulated in the cell to which the gene is native. For example, an endogenous gene of a cell can be regulated by an exogenous regulatory sequence inserted into the genome of the cell at location sufficient to at least enhance expressed of the gene in the cell. The regulatory sequence may be designed to integrate into the genome via homologous recombination, as disclosed in U.S. Patent Nos. 5,641,670 and 5,733,761, the disclosures of which are herein incorporated by reference, or may be designed to integrate into the genome via non-homologous recombination, as described in WO 99/15650, the disclosure of which is herein incorporated by reference. As such, also encompassed in the subject invention is the production of the subject proteins without manipulation of the encoding nucleic acid itself, but instead through integration of a regulatory sequence into the genome of cell that already includes a gene encoding the desired protein, as described in the above incorporated patent documents.

Also provided are homologs of the subject nucleic acids. Homologs are identified by any of a number of methods. A fragment of the provided cDNA may be used as a hybridization probe against a cDNA library from the target organism of interest, where low stringency conditions are used. The probe may be a large fragment, or one or more short degenerate primers. Nucleic acids having sequence similarity are detected by hybridization under low stringency conditions, for example, at 50°C and 6×SSC (0.9 M sodium chloride/0.09 M sodium citrate) and remain bound when subjected to washing at

55°C in 1×SSC (0.15 M sodium chloride/0.015 M sodium citrate). Sequence identity may be determined by hybridization under stringent conditions, for example, at 50°C or higher and 0.1×SSC (15 mM sodium chloride/1.5 mM sodium citrate). Nucleic acids having a region of substantial identity to the provided sequences, e.g. allelic variants, genetically altered versions of the gene, etc., bind to the provided sequences under stringent hybridization conditions. By using probes, particularly labeled probes of DNA sequences, one can isolate homologous or related genes.

Also of interest are promoter elements of the subject genomic sequences, where the sequence of the 5' flanking region may be utilized for promoter elements, including enhancer binding sites, e.g., that provide for regulation of expression in cells/tissues where the subject proteins gene are expressed.

Also provided are small DNA fragments of the subject nucleic acids, which fragments are useful as primers for PCR, hybridization screening probes, etc. Larger DNA fragments, i.e., greater than 100 nt are useful for production of the encoded polypeptide, as described in the previous section. For use in geometric amplification reactions, such as geometric PCR, a pair of primers will be used. The exact composition of the primer sequences is not critical to the invention, but for most applications the primers will hybridize to the subject sequence under stringent conditions, as known in the art. It is preferable to choose a pair of primers that will generate an amplification product of at least about 50 nt, preferably at least about 100 nt. Algorithms for the selection of primer sequences are generally known, and are available in commercial software packages. Amplification primers hybridize to complementary strands of DNA, and will prime towards each other.

The DNA may also be used to identify expression of the gene in a biological specimen. The manner in which one probes cells for the presence of particular nucleotide sequences, as genomic DNA or RNA, is well established in the literature. Briefly, DNA or mRNA is isolated from a cell sample. The mRNA may be amplified by RT-PCR, using reverse transcriptase to form a complementary DNA strand, followed by polymerase chain reaction amplification using primers specific for the subject DNA sequences.

Alternatively, the mRNA sample is separated by gel electrophoresis, transferred to a suitable support, e.g. nitrocellulose, nylon, etc., and then probed with a fragment of the subject DNA as a probe. Other techniques, such as oligonucleotide ligation assays, *in situ* hybridizations, and hybridization to DNA probes arrayed on a solid chip may also find use. Detection of mRNA hybridizing to the subject sequence is indicative of Anthozoan protein gene expression in the sample.

The subject nucleic acids, including flanking promoter regions and coding regions, may be mutated in various ways known in the art to generate targeted changes in promoter strength, sequence of the encoded protein, properties of the encoded protein, including fluorescent properties of the encoded protein, etc. The DNA sequence or protein

product of such a mutation will usually be substantially similar to the sequences provided herein, e.g. will differ by at least one nucleotide or amino acid, respectively, and may differ by at least two but not more than about ten nucleotides or amino acids. The sequence changes may be substitutions, insertions, deletions, or a combination thereof. Deletions may further include larger changes, such as deletions of a domain or exon, e.g. of stretches of 10, 20, 50, 75, 100, 150 or more aa residues. Techniques for *in vitro* mutagenesis of cloned genes are known. Examples of protocols for site specific mutagenesis may be found in Gustin *et al.* (1993), *Biotechniques* 14:22; Barany (1985), *Gene* 37:111-23; Colicelli *et al.* (1985), *Mol. Gen. Genet.* 199:537-9; and Prentki *et al.* (1984), *Gene* 29:303-13. Methods for site specific mutagenesis can be found in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, CSH Press 1989, pp. 15.3-15.108; Weiner *et al.* (1993), *Gene* 126:35-41; Sayers *et al.* (1992), *Biotechniques* 13:592-6; Jones and Winistorfer (1992), *Biotechniques* 12:528-30; Barton *et al.* (1990), *Nucleic Acids Res* 18:7349-55; Marotti and Tomich (1989), *Gene Anal. Tech.* 6:67-70; and Zhu (1989), *Anal Biochem* 177:120-4. Such mutated nucleic acid derivatives may be used to study structure-function relationships of a particular chromo/ fluorescent protein, or to alter properties of the protein that affect its function or regulation.

Of particular interest in many embodiments is the following specific mutation protocol, which protocol finds use in mutating chromoproteins (e.g., colored proteins that have little if any fluorescence) into fluorescent mutants. In this protocol, the sequence of the candidate protein is aligned with the amino acid sequence of *Aequorea victoria* wild type GFP, according to the protocol reported in Matz *et al.*, "Fluorescent proteins from non-bioluminescent Anthozoa species," *Nature Biotechnology* (October 1999) 17: 969 – 973. Residue 148 of the aligned chromoprotein is identified and then changed to Ser, e.g., by site directed mutagenesis, which results in the production of a fluorescent mutant of the wild type chromoprotein. See e.g., NFP-7 described below, which wild type protein is a chromoprotein that is mutated into a fluorescent protein by substitution of Ser for the native Ala residue at position 148.

Also of interest are humanized versions of the subject nucleic acids. As used herein, the term "humanized" refers to changes made to the a nucleic acid sequence to optimize the codons for expression of the protein in human cells (Yang *et al.*, *Nucleic Acids Research* 24 (1996), 4592-4593). See also U.S. Patent No. 5,795,737 which describes humanization of proteins, the disclosure of which is herein incorporated by reference.

PROTEIN/POLYPEPTIDE COMPOSITIONS

Also provided by the subject invention are chromo- and/or fluorescent proteins and mutants thereof, as well as polypeptide compositions related thereto. As the subject proteins are chromoproteins, they are colored proteins, which may be fluorescent, low or non- fluorescent. As used herein, the terms chromoprotein and fluorescent protein do not include luciferases, such as Renilla luciferase, and refer to any protein that is pigmented or colored and/or fluoresces when irradiated with light, e.g., white light or light of a specific wavelength (or narrow band of wavelengths such as an excitation wavelength). The term polypeptide composition as used herein refers to both the full-length protein, as well as portions or fragments thereof. Also included in this term are variations of the naturally occurring protein, where such variations are homologous or substantially similar to the naturally occurring protein, and mutants of the naturally occurring proteins, as described in greater detail below. The subject polypeptides are present in other than their natural environment.

In many embodiments, the subject proteins have an absorbance maximum ranging from about 300 to 700, usually from about 350 to 650 and more usually from about 400 to 600 nm. Where the subject proteins are fluorescent proteins, by which is meant that they can be excited at one wavelength of light following which they will emit light at another wavelength, the excitation spectra of the subject proteins typically ranges from about 300 to 700, usually from about 350 to 650 and more usually from about 400 to 600 nm while the emission spectra of the subject proteins typically ranges from about 400 to 800, usually from about 425 to 775 and more usually from about 450 to 750 nm. The subject proteins generally have a maximum extinction coefficient that ranges from about 10,000 to 50,000 and usually from about 15,000 to 45,000. The subject proteins typically range in length from about 150 to 300 and usually from about 200 to 300 amino acid residues, and generally have a molecular weight ranging from about 15 to 35 kDa, usually from about 17.5 to 32.5 kDa.

In certain embodiments, the subject proteins are bright, where by bright is meant that the chromoproteins and their fluorescent mutants can be detected by common methods (e.g., visual screening, spectrophotometry, spectrofluorometry, fluorescent microscopy, by FACS machines, etc.) Fluorescence brightness of particular fluorescent proteins is determined by its quantum yield multiplied by maximal extinction coefficient. Brightness of a chromoproteins may be expressed by its maximal extinction coefficient.

In certain embodiments, the subject proteins fold rapidly following expression in the host cell. By rapidly folding is meant that the proteins achieve their tertiary structure that gives rise to their chromo- or fluorescent quality in a short period of time. In these embodiments, the proteins fold in a period of time that generally does not exceed about 3 days, usually does not exceed about 2 days and more usually does not exceed about 1 day.

Specific proteins of interest are chromo/fluoroproteins (and mutants thereof) from the following specific Anthozoan species: *Anemonia majano*, *Clavularia* sp., *Zoanthus* sp., *Zoanthus* sp., *Discosoma striata*, *Discosoma* sp. "red", *Anemonia sulcata*, *Discosoma* sp "green", *Discosoma* sp."magenta." Each of these particular types of polypeptide compositions of interest is now discussed in greater detail individually.

Anemonia majano (NFP-1; AmCyan)

In many embodiments, the proteins have an absorbance maximum ranging from about 250 to 650, usually from about 400 to 500 and more usually from about 440 to 480 nm while the emission maximum typically ranges from about 270 to 670, usually from about 420 to 520 and more usually from about 460 to 500 nm. The subject proteins typically range in length from about 200 to 250, usually from about 210 to 240 amino acid residues, and generally have a molecular weight ranging from about 20 to 30, usually from about 22.50 to 27.50 kDa. Of particular interest in many embodiments is amFP486 (NFP-1), which has an amino acid sequence as shown in SEQ ID NO:02. Also of interest are mutants of this sequence, where specific mutants of interest include, but are not limited to: Mut15, Mut32, and FP1-NA (a non-aggregating mutant), where these specific mutants are further described in the experimental section, infra.

Clavularia sp. (NFP-2)

In certain embodiments, the proteins have an absorbance maximum that typically ranges from about 250 to 650, usually from about 400 to 500 and more usually from about 440 to 480 nm and an emission maximum that typically ranges from about 270 to 670, usually from about 420 to 520 and more usually from about 460 to 500 nm, where the subject proteins typically range in length from about 225 to 300, usually from about 250 to 275 amino acid residues, and generally have a molecular weight ranging from about 25 to 35, usually from about 27.50 to 32.50 kDa. Of particular interest is the cFP484 protein having the sequence shown in SEQ ID NO:04. Specific mutants of interest include, but are not limited to: $\Delta 19$ cFP484 and $\Delta 38$ cFP484, where these specific mutants are further described in the experimental section, infra.

Zoanthus sp I. (NFP-3; ZsGreen)

In many embodiments, the proteins have an absorbance maximum that typically ranges from about 300 to 700, usually from about 450 to 550 and more usually from about 480 to 510 nm and an emission maximum that typically ranges from about 320 to 720, usually from about 470 to 570 and more usually from about 500 to 530 nm. The subject

proteins typically range in length from about 200 to 250, usually from about 220 to 240 amino acid residues, and generally have a molecular weight ranging from about 20 to 30, usually from about 22.50 to 27.50 kDa. Of particular interest is the protein zFP506 (NFP-3) which has an amino acid sequence as shown in SEQ ID NO:06. Specific mutants of interest include, but are not limited to: N66M; FP-3NA (a non-aggregating mutant); yellow; yellow bright, etc., where these specific mutants are further described in the experimental section, infra.

Zoanthus sp. II (NFP-4; ZsYellow)

In many embodiments, the proteins have an excitation maximum that typically ranges from about 300 to 650, usually from about 475 to 575 and more usually from about 500 to 550 nm and an emission maximum that typically ranges from about 310 to 660, usually from about 485 to 585 and more usually from about 510 to 560 nm. The subject proteins typically range in length from about 200 to 250, usually from about 220 to 240 amino acid residues, and generally have a molecular weight ranging from about 20 to 30, usually from about 22.50 to 27.50 kDa. Specific mutants of interest include, but are not limited to: M129V; NFP4-NA (a non-aggregating mutant); green; etc., where these specific mutants are further described in the experimental section, infra.

Discosoma striata (NFP-5)

In many embodiments, the proteins have an excitation maximum that typically ranges from about 240 to 640, usually from about 500 to 600 and more usually from about 530 to 560 nm and an emission maximum that typically ranges from about 280 to 680, usually from about 540 to 640 and more usually from about 570 to 600 nm. The subject proteins typically range in length from about 200 to 250, usually from about 220 to 240 amino acid residues, and generally have a molecular weight ranging from about 20 to 30, usually from about 22.50 to 27.50 kDa. Of particular interest in many embodiments is the protein dsFP483 (NFP-5) which has an amino acid sequence as shown in SEQ ID NO:10, as well as mutants thereof.

Discosoma sp. "red" (NFP-6; DsRed; DsRed2)

In many embodiments, the proteins have an absorbance maximum that typically ranges from about 250 to 750, usually from about 500 to 600 and more usually from about 540 to 580 nm and have an emission maximum that typically ranges from about 275 to 775, usually from about 525 to 625 and more usually from about 565 to 605 nm. The

subject proteins typically range in length from about 200 to 250, usually from about 220 to 240 amino acid residues, and generally have a molecular weight ranging from about 20 to 30, usually from about 22.50 to 27.50 kDa. Of particular interest is the drFP583 (NFP-6) protein that has an amino acid sequence as shown in SEQ ID NO:12. Specific mutants of interest include, but are not limited to: E5, E5-NA (a non-aggregating mutant), E8, E5up, E5down, E57, FP6-NA (a non-aggregating mutant), AG4, AG45, E83, 6/9 Q, 6/9 Q-NA, 6/92G, etc., where these specific mutants are further described in the experimental section, *infra*.

10 *Anemonia sulcata* (NFP-7; AsRed)

In many embodiments, the proteins have an absorbance maximum that typically ranges from about 370 to 770, usually from about 520 to 620 and more usually from about 560 to 580 nm and an emission maximum that typically ranges from about 395 to 795, usually from about 545 to 645 and more usually from about 585 to 605 nm. The subject proteins typically range in length from about 200 to 250, usually from about 220 to 240 amino acid residues, and generally have a molecular weight ranging from about 20 to 30, usually from about 22.50 to 27.50 kDa. Of particular interest is the asFP595(asFP600) (NFP-7) protein that has an amino acid sequence as shown in SEQ ID NO:14. Specific mutants of interest include, but are not limited to: Mut1; Mut35-5/Mut1; FP7-NA (a non-aggregating mutant), etc., where these specific mutants are further described in the experimental section, *infra*.

25 *Discosoma* sp "green" (NFP-8)

In many embodiments, the proteins have an absorbance maximum that typically ranges from about 300 to 700, usually from about 450 to 650 and more usually from about 490 to 510 nm and an emission maximum that typically ranges from about 310 to 710, usually from about 460 to 660 and more usually from about 500 to 520 nm. The subject proteins typically range in length from about 200 to 250, usually from about 220 to 240 amino acid residues, and generally have a molecular weight ranging from about 20 to 30, usually from about 22.50 to 27.50 kDa. Of particular interest is the dgFP512 protein (NFP-8) protein that has an amino acid sequence as shown in SEQ ID NO:16, as well as mutants thereof.

35 *Discosoma* sp. "magenta" (NFP-9)

In many embodiments, the proteins have an absorbance maximum that typically ranges from about 375 to 775, usually from about 525 to 625 and more usually from about

560 to 590 nm and an emission maximum that typically ranges from about 395 to 795, usually from about 545 to 645 and more usually from about 580 to 610 nm. The subject proteins typically range in length from about 200 to 250, usually from about 220 to 240 amino acid residues, and generally have a molecular weight ranging from about 20 to 30, usually from about 22.50 to 27.50 kDa. Of particular interest is the dmFP592 (NFP-9) protein that has an amino acid sequence as shown in SEQ ID NO:18, as well as mutants thereof.

Homologs or proteins (or fragments thereof) that vary in sequence from the above provided specific amino acid sequences of the subject invention, i.e., SEQ ID NOS: 02; 04; 06; 08; 10; 12; 14; 16 or 18, are also provided. By homolog is meant a protein having at least about 10%, usually at least about 20 % and more usually at least about 30 %, and in many embodiments at least about 35 %, usually at least about 40% and more usually at least about 60 % amino acid sequence identity to the protein of the subject invention, as determined using MegAlign, DNASTAR (1998) clustal algorithm as described in D. G. Higgins and P.M. Sharp, "Fast and Sensitive multiple Sequence Alignments on a Microcomputer," (1989) CABIOS, 5: 151-153. (Parameters used are ktuple 1, gap penalty 3, window, 5 and diagonals saved 5). In many embodiments, homologues of interest have much higher sequence identity, e.g., 65%, 70%, 75%, 80%, 85%, 90% or higher.

Also provided are proteins that are substantially identical to the wild type protein, where by substantially identical is meant that the protein has an amino acid sequence identity to the sequence of wild type protein of at least about 60%, usually at least about 65% and more usually at least about 70 %, where in some instances the identity may be much higher, e.g., 75%, 80%, 85%, 90%, 95% or higher.

In many embodiments, the subject homologues have structural features found in the above provided specific sequences, where such structural features include the β -can fold.

Proteins which are mutants of the above-described naturally occurring proteins are also provided. Mutants may retain biological properties of the wild-type (e.g., naturally occurring) proteins, or may have biological properties which differ from the wild-type proteins. The term "biological property" of the subject proteins includes, but is not limited to, spectral properties, such as absorbance maximum, emission maximum, maximum extinction coefficient, brightness (e.g., as compared to the wild-type protein or another reference protein such as green fluorescent protein from *A. victoria*), and the like; *in vivo* and/or *in vitro* stability (e.g., half-life); etc. Mutants include single amino acid changes, deletions of one or more amino acids, N-terminal truncations, C-terminal truncations, insertions, etc.

1 Mutants can be generated using standard techniques of molecular biology, e.g.,
2 random mutagenesis, and targeted mutagenesis. Several mutants are described herein.
3 Given the guidance provided in the Examples, and using standard techniques, those
4 skilled in the art can readily generate a wide variety of additional mutants and test whether
5 a biological property has been altered. For example, fluorescence intensity can be
6 measured using a spectrophotometer at various excitation wavelengths.

7 Those proteins of the subject invention that are naturally occurring proteins are
8 present in a non-naturally occurring environment, e.g., are separated from their naturally
9 occurring environment. In certain embodiments, the subject proteins are present in a
10 composition that is enriched for the subject protein as compared to its naturally occurring
11 environment. For example, purified protein is provided, where by purified is meant that the
12 protein is present in a composition that is substantially free of non- chromo/fluoroprotein
13 proteins of interest, where by substantially free is meant that less than 90 %, usually less
14 than 60 % and more usually less than 50 % of the composition is made up of non-
15 chromoproteins or mutants thereof of interest. The proteins of the subject invention may
16 also be present as an isolate, by which is meant that the protein is substantially free of
17 other proteins and other naturally occurring biologic molecules, such as oligosaccharides,
18 polynucleotides and fragments thereof, and the like, where the term "substantially free" in
19 this instance means that less than 70 %, usually less than 60% and more usually less
20 than 50 % of the composition containing the isolated protein is some other naturally
21 occurring biological molecule. In certain embodiments, the proteins are present in
22 substantially pure form, where by "substantially pure form" is meant at least 95%, usually
23 at least 97% and more usually at least 99% pure.

24 In addition to the naturally occurring proteins, polypeptides that vary from the
25 naturally occurring proteins, e.g., the mutant proteins described above, are also provided.
26 Generally such polypeptides include an amino acid sequence encoded by an open
27 reading frame (ORF) of the gene encoding the subject wild type protein, including the full
28 length protein and fragments thereof, particularly biologically active fragments and/or
29 fragments corresponding to functional domains, and the like; and including fusions of the
30 subject polypeptides to other proteins or parts thereof. Fragments of interest will typically
31 be at least about 10 aa in length, usually at least about 50 aa in length, and may be as
32 long as 300 aa in length or longer, but will usually not exceed about 1000 aa in length,
33 where the fragment will have a stretch of amino acids that is identical to the subject
34 protein of at least about 10 aa, and usually at least about 15 aa, and in many
35 embodiments at least about 50 aa in length. In some embodiments, the subject
36 polypeptides are about 25 aa, about 50 aa, about 75 aa, about 100 aa, about 125 aa,
37 about 150 aa, about 200 aa, about 210 aa, about 220 aa, about 230 aa, or about 240 aa
38 in length, up to the entire protein. In some embodiments, a protein fragment retains all or
39 substantially all of a biological property of the wild-type protein.

The subject proteins and polypeptides may be obtained from naturally occurring sources or synthetically produced. For example, wild type proteins may be derived from biological sources which express the proteins, e.g., non-bioluminescent Cnidarian, e.g., Anthozoan, species, such as the specific ones listed above. The subject proteins may also be derived from synthetic means, e.g., by expressing a recombinant gene or nucleic acid coding sequence encoding the protein of interest in a suitable host, as described above. Any convenient protein purification procedures may be employed, where suitable protein purification methodologies are described in Guide to Protein Purification, (Deuthser ed.) (Academic Press, 1990). For example, a lysate may prepared from the original source and purified using HPLC, exclusion chromatography, gel electrophoresis, affinity chromatography, and the like.

ANTIBODY COMPOSITIONS

Also provided are antibodies that specifically bind to the subject fluorescent proteins. Suitable antibodies are obtained by immunizing a host animal with peptides comprising all or a portion of the subject protein. Suitable host animals include mouse, rat, sheep, goat, hamster, rabbit, etc. The origin of the protein immunogen will generally be a Cnidarian species, specifically a non-bioluminescent Cnidarian species, such as an Anthozoan species or a non-Petalucean Anthozoan species. The host animal will generally be a different species than the immunogen, e.g., mice, etc.

The immunogen may comprise the complete protein, or fragments and derivatives thereof. Preferred immunogens comprise all or a part of the protein, where these residues contain the post-translation modifications found on the native target protein.

Immunogens are produced in a variety of ways known in the art, e.g., expression of cloned genes using conventional recombinant methods, isolation from Anthozoan species of origin, etc.

For preparation of polyclonal antibodies, the first step is immunization of the host animal with the target protein, where the target protein will preferably be in substantially pure form, comprising less than about 1% contaminant. The immunogen may comprise the complete target protein, fragments or derivatives thereof. To increase the immune response of the host animal, the target protein may be combined with an adjuvant, where suitable adjuvants include alum, dextran, sulfate, large polymeric anions, oil & water emulsions, e.g. Freund's adjuvant, Freund's complete adjuvant, and the like. The target protein may also be conjugated to synthetic carrier proteins or synthetic antigens. A variety of hosts may be immunized to produce the polyclonal antibodies. Such hosts include rabbits, guinea pigs, rodents, e.g. mice, rats, sheep, goats, and the like. The target protein is administered to the host, usually intradermally, with an initial dosage followed by one or more, usually at least two, additional booster dosages. Following

immunization, the blood from the host will be collected, followed by separation of the serum from the blood cells. The Ig present in the resultant antiserum may be further fractionated using known methods, such as ammonium salt fractionation, DEAE chromatography, and the like.

Monoclonal antibodies are produced by conventional techniques. Generally, the spleen and/or lymph nodes of an immunized host animal provide a source of plasma cells. The plasma cells are immortalized by fusion with myeloma cells to produce hybridoma cells. Culture supernatant from individual hybridomas is screened using standard techniques to identify those producing antibodies with the desired specificity. Suitable animals for production of monoclonal antibodies to the human protein include mouse, rat, hamster, etc. To raise antibodies against the mouse protein, the animal will generally be a hamster, guinea pig, rabbit, etc. The antibody may be purified from the hybridoma cell supernatants or ascites fluid by conventional techniques, e.g. affinity chromatography using protein bound to an insoluble support, protein A sepharose, etc.

The antibody may be produced as a single chain, instead of the normal multimeric structure. Single chain antibodies are described in Jost *et al.* (1994) J.B.C. 269:26267-73, and others. DNA sequences encoding the variable region of the heavy chain and the variable region of the light chain are ligated to a spacer encoding at least about 4 amino acids of small neutral amino acids, including glycine and/or serine. The protein encoded by this fusion allows assembly of a functional variable region that retains the specificity and affinity of the original antibody.

Also of interest in certain embodiments are humanized antibodies. Methods of humanizing antibodies are known in the art. The humanized antibody may be the product of an animal having transgenic human immunoglobulin constant region genes (see for example International Patent Applications WO 90/10077 and WO 90/04036). Alternatively, the antibody of interest may be engineered by recombinant DNA techniques to substitute the CH1, CH2, CH3, hinge domains, and/or the framework domain with the corresponding human sequence (see WO 92/02190).

The use of Ig cDNA for construction of chimeric immunoglobulin genes is known in the art (Liu *et al.* (1987) P.N.A.S. 84:3439 and (1987) J. Immunol. 139:3521). mRNA is isolated from a hybridoma or other cell producing the antibody and used to produce cDNA. The cDNA of interest may be amplified by the polymerase chain reaction using specific primers (U.S. Patent nos. 4,683,195 and 4,683,202). Alternatively, a library is made and screened to isolate the sequence of interest. The DNA sequence encoding the variable region of the antibody is then fused to human constant region sequences. The sequences of human constant regions genes may be found in Kabat *et al.* (1991) Sequences of Proteins of Immunological Interest, N.I.H. publication no. 91-3242. Human C region genes are readily available from known clones. The choice of isotype will be guided by the desired effector functions, such as complement fixation, or activity in

antibody-dependent cellular cytotoxicity. Preferred isotypes are IgG1, IgG3 and IgG4. Either of the human light chain constant regions, kappa or lambda, may be used. The chimeric, humanized antibody is then expressed by conventional methods.

Antibody fragments, such as Fv, F(ab')₂ and Fab may be prepared by cleavage of the intact protein, *e.g.* by protease or chemical cleavage. Alternatively, a truncated gene is designed. For example, a chimeric gene encoding a portion of the F(ab')₂ fragment would include DNA sequences encoding the CH1 domain and hinge region of the H chain, followed by a translational stop codon to yield the truncated molecule.

Consensus sequences of H and L J regions may be used to design oligonucleotides for use as primers to introduce useful restriction sites into the J region for subsequent linkage of V region segments to human C region segments. C region cDNA can be modified by site directed mutagenesis to place a restriction site at the analogous position in the human sequence.

Expression vectors include plasmids, retroviruses, YACs, EBV derived episomes, and the like. A convenient vector is one that encodes a functionally complete human CH or CL immunoglobulin sequence, with appropriate restriction sites engineered so that any VH or VL sequence can be easily inserted and expressed. In such vectors, splicing usually occurs between the splice donor site in the inserted J region and the splice acceptor site preceding the human C region, and also at the splice regions that occur within the human CH exons. Polyadenylation and transcription termination occur at native chromosomal sites downstream of the coding regions. The resulting chimeric antibody may be joined to any strong promoter, including retroviral LTRs, *e.g.* SV-40 early promoter, (Okayama *et al.* (1983) Mol. Cell. Bio. 3:280), Rous sarcoma virus LTR (Gorman *et al.* (1982) P.N.A.S. 79:6777), and moloney murine leukemia virus LTR (Grosschedl *et al.* (1985) Cell 41:885); native Ig promoters, *etc.*

TRANSGENICS

The subject nucleic acids can be used to generate transgenic, non-human plants or animals or site specific gene modifications in cell lines. Transgenic cells of the subject invention include on or more nucleic acids according to the subject invention present as a transgene, where included within this definition are the parent cells transformed to include the transgene and the progeny thereof. In many embodiments, the transgenic cells are cells that do not normally harbor or contain a nucleic acid according to the subject invention. In those embodiments where the transgenic cells do naturally contain the subject nucleic acids, the nucleic acid will be present in the cell in a position other than its natural location, *i.e.* integrated into the genomic material of the cell at a non-natural location. Transgenic animals may be made through homologous recombination, where the endogenous locus is altered. Alternatively, a nucleic acid construct is randomly

integrated into the genome. Vectors for stable integration include plasmids, retroviruses and other animal viruses, YACs, and the like.

Transgenic organisms of the subject invention include cells and multicellular organisms, e.g., plants and animals, that are endogenous knockouts in which expression of the endogenous gene is at least reduced if not eliminated. Transgenic organisms of interest also include cells and multicellular organisms, e.g., plants and animals, in which the protein or variants thereof is expressed in cells or tissues where it is not normally expressed and/or at levels not normally present in such cells or tissues.

DNA constructs for homologous recombination will comprise at least a portion of the gene of the subject invention, wherein the gene has the desired genetic modification(s), and includes regions of homology to the target locus. DNA constructs for random integration need not include regions of homology to mediate recombination. Conveniently, markers for positive and negative selection are included. Methods for generating cells having targeted gene modifications through homologous recombination are known in the art. For various techniques for transfecting mammalian cells, see Keown *et al.* (1990), *Meth. Enzymol.* 185:527-537.

For embryonic stem (ES) cells, an ES cell line may be employed, or embryonic cells may be obtained freshly from a host, e.g. mouse, rat, guinea pig, etc. Such cells are grown on an appropriate fibroblast-feeder layer or grown in the presence of leukemia inhibiting factor (LIF). When ES or embryonic cells have been transformed, they may be used to produce transgenic animals. After transformation, the cells are plated onto a feeder layer in an appropriate medium. Cells containing the construct may be detected by employing a selective medium. After sufficient time for colonies to grow, they are picked and analyzed for the occurrence of homologous recombination or integration of the construct. Those colonies that are positive may then be used for embryo manipulation and blastocyst injection. Blastocysts are obtained from 4 to 6 week old superovulated females. The ES cells are trypsinized, and the modified cells are injected into the blastocoel of the blastocyst. After injection, the blastocysts are returned to each uterine horn of pseudopregnant females. Females are then allowed to go to term and the resulting offspring screened for the construct. By providing for a different phenotype of the blastocyst and the genetically modified cells, chimeric progeny can be readily detected.

The chimeric animals are screened for the presence of the modified gene and males and females having the modification are mated to produce homozygous progeny. If the gene alterations cause lethality at some point in development, tissues or organs can be maintained as allogeneic or congenic grafts or transplants, or in *in vitro* culture. The transgenic animals may be any non-human mammal, such as laboratory animals, domestic animals, etc. The transgenic animals may be used in functional studies, drug

screening, etc. Representative examples of the use of transgenic animals include those described infra.

Transgenic plants may be produced in a similar manner. Methods of preparing transgenic plant cells and plants are described in U.S. Pat. Nos. 5,767,367; 5,750,870; 5,739,409; 5,689,049; 5,689,045; 5,674,731; 5,656,466; 5,633,155; 5,629,470 ; 5,595,896; 5,576,198; 5,538,879; 5,484,956; the disclosures of which are herein incorporated by reference. Methods of producing transgenic plants are also reviewed in Plant Biochemistry and Molecular Biology (eds Lea & Leegood, John Wiley & Sons)(1993) pp 275-295. In brief, a suitable plant cell or tissue is harvested, depending on the nature of the plant species. As such, in certain instances, protoplasts will be isolated, where such protoplasts may be isolated from a variety of different plant tissues, e.g. leaf, hypocotyl, root, etc. For protoplast isolation, the harvested cells are incubated in the presence of cellulases in order to remove the cell wall, where the exact incubation conditions vary depending on the type of plant and/or tissue from which the cell is derived. The resultant protoplasts are then separated from the resultant cellular debris by sieving and centrifugation. Instead of using protoplasts, embryogenic explants comprising somatic cells may be used for preparation of the transgenic host. Following cell or tissue harvesting, exogenous DNA of interest is introduced into the plant cells, where a variety of different techniques are available for such introduction. With isolated protoplasts, the opportunity arise for introduction via DNA-mediated gene transfer protocols, including: incubation of the protoplasts with naked DNA, e.g. plasmids, comprising the exogenous coding sequence of interest in the presence of polyvalent cations, e.g. PEG or PLO; and electroporation of the protoplasts in the presence of naked DNA comprising the exogenous sequence of interest. Protoplasts that have successfully taken up the exogenous DNA are then selected, grown into a callus, and ultimately into a transgenic plant through contact with the appropriate amounts and ratios of stimulatory factors, e.g. auxins and cytokinins. With embryogenic explants, a convenient method of introducing the exogenous DNA in the target somatic cells is through the use of particle acceleration or "gene-gun" protocols. The resultant explants are then allowed to grow into chimera plants, cross-bred and transgenic progeny are obtained. Instead of the naked DNA approaches described above, another convenient method of producing transgenic plants is *Agrobacterium* mediated transformation. With *Agrobacterium* mediated transformation, co-integrative or binary vectors comprising the exogenous DNA are prepared and then introduced into an appropriate *Agrobacterium* strain, e.g. *A. tumefaciens*. The resultant bacteria are then incubated with prepared protoplasts or tissue explants, e.g. leaf disks, and a callus is produced. The callus is then grown under selective conditions, selected and subjected to growth media to induce root and shoot growth to ultimately produce a transgenic plant.

UTILITY

The subject chromoproteins and fluorescent mutants thereof find use in a variety of different applications, where the applications necessarily differ depending on whether the protein is a chromoprotein or a fluorescent protein. Representative uses for each of these types of proteins will be described below, where the follow described uses are merely representative and are in no way meant to limit the use of the subject proteins to those described below.

Chromoproteins

The subject chromoproteins of the present invention find use in a variety of different applications. One application of interest is the use of the subject proteins as coloring agents which are capable of imparting color or pigment to a particular composition of matter. Of particular interest in certain embodiments are non-toxic chromoproteins. The subject chromoproteins may be incorporated into a variety of different compositions of matter, where representative compositions of matter include: food compositions, pharmaceuticals, cosmetics, living organisms, e.g., animals and plants, and the like. Where used as a coloring agent or pigment, a sufficient amount of the chromoprotein is incorporated into the composition of matter to impart the desired color or pigment thereto. The chromoprotein may be incorporated into the composition of matter using any convenient protocol, where the particular protocol employed will necessarily depend, at least in part, on the nature of the composition of matter to be colored.

Protocols that may be employed include, but are not limited to: blending, diffusion, friction, spraying, injection, tattooing, and the like.

The chromoproteins may also find use as labels in analyte detection assays, e.g., assays for biological analytes of interest. For example, the chromoproteins may be incorporated into adducts with analyte specific antibodies or binding fragments thereof and subsequently employed in immunoassays for analytes of interest in a complex sample, as described in U.S. Patent No. 4,302,536; the disclosure of which is herein incorporated by reference. Instead of antibodies or binding fragments thereof, the subject chromoproteins or chromogenic fragments thereof may be conjugated to ligands that specifically bind to an analyte of interest, or other moieties, growth factors, hormones, and the like; as is readily apparent to those of skill in the art.

In yet other embodiments, the subject chromoproteins may be used as selectable markers in recombinant DNA applications, e.g., the production of transgenic cells and organisms, as described above. As such, one can engineer a particular transgenic production protocol to employ expression of the subject chromoproteins as a selectable

marker, either for a successful or unsuccessful protocol. Thus, appearance of the color of the subject chromoprotein in the phenotype of the transgenic organism produced by a particular process can be used to indicate that the particular organism successfully harbors the transgene of interest, often integrated in a manner that provides for expression of the transgene in the organism. When used a selectable marker, a nucleic acid encoding for the subject chromoprotein can be employed in the transgenic generation process, where this process is described in greater detail supra. Particular transgenic organisms of interest where the subject proteins may be employed as selectable markers include transgenic plants, animals, bacteria, fungi, and the like.

In yet other embodiments, the chromoproteins (and fluorescent proteins) of the subject invention find use in sunscreens, as selective filters, etc., in a manner similar to the uses of the proteins described in WO 00/46233.

Fluorescent Proteins

The subject fluorescent proteins of the present invention (as well as other components of the subject invention described above) find use in a variety of different applications, where such applications include, but are not limited to, the following. The first application of interest is the use of the subject proteins in fluorescence resonance energy transfer (FRET) applications. In these applications, the subject proteins serve as donor and/or acceptors in combination with a second fluorescent protein or dye, e.g., a fluorescent protein as described in Matz et al., Nature Biotechnology (October 1999) 17:969-973, a green fluorescent protein from *Aequoria victoria* or fluorescent mutant thereof, e.g., as described in U.S. Patent No. 6,066,476; 6,020,192; 5,985,577; 5,976,796; 5,968,750; 5,968,738; 5,958,713; 5,919,445; 5,874,304, the disclosures of which are herein incorporated by reference, other fluorescent dyes, e.g., coumarin and its derivatives, e.g. 7-amino-4-methylcoumarin, aminocoumarin, bodipy dyes, such as Bodipy FL, cascade blue, fluorescein and its derivatives, e.g. fluorescein isothiocyanate, Oregon green, rhodamine dyes, e.g. texas red, tetramethylrhodamine, eosins and erythrosins, cyanine dyes, e.g. Cy3 and Cy5, macrocyclic chelates of lanthanide ions, e.g. quantum dye, etc., chemiluminescent dyes, e.g., luciferases, including those described in U.S. Patent Nos. 5,843,746; 5,700,673; 5,674,713; 5,618,722; 5,418,155; 5,330,906; 5,229,285; 5,221,623; 5,182,202; the disclosures of which are herein incorporated by reference. Specific examples of where FRET assays employing the subject fluorescent proteins may be used include, but are not limited to: the detection of protein-protein interactions, e.g., mammalian two-hybrid system, transcription factor dimerization, membrane protein multimerization, multiprotein complex formation, etc., as a biosensor for a number of different events, where a peptide or protein covalently links a FRET fluorescent combination including the subject fluorescent proteins and the linking peptide or protein is,

e.g., a protease specific substrate, e.g., for caspase mediated cleavage, a linker that undergoes conformational change upon receiving a signal which increases or decreases FRET, e.g., PKA regulatory domain (cAMP-sensor), phosphorylation, e.g., where there is a phosphorylation site in the linker or the linker has binding specificity to

5 phosphorylated/dephosphorylated domain of another protein, or the linker has Ca^{2+} binding domain. Representative fluorescence resonance energy transfer or FRET applications in which the subject proteins find use include, but are not limited to, those described in: U.S. Patent Nos. 6,008,373; 5,998,146; 5,981,200; 5,945,526; 5,945,283; 5,911,952; 5,869,255; 5,866,336; 5,863,727; 5,728,528; 5,707,804; 5,688,648; 10 5,439,797; the disclosures of which are herein incorporated by reference.

The subject fluorescent proteins also find use as biosensors in prokaryotic and eukaryotic cells, e.g. as Ca^{2+} ion indicator; as pH indicator, as phosphorylation indicator, as an indicator of other ions, e.g., magnesium, sodium, potassium, chloride and halides. For example, for detection of Ca ion, proteins containing an EF-hand motif are known to translocate from the cytosol to membranes upon Ca^{2+} binding. These proteins contain a myristoyl group that is buried within the molecule by hydrophobic interactions with other regions of the protein. Binding of Ca^{2+} induces a conformational change exposing the myristoyl group which then is available for the insertion into the lipid bilayer (called a " Ca^{2+} -myristoyl switch"). Fusion of such a EF-hand containing protein to Fluorescent Proteins 15 (FP) could make it an indicator of intracellular Ca^{2+} by monitoring the translocation from the cytosol to the plasma membrane by confocal microscopy. EF-hand proteins suitable for use in this system include, but are not limited to: recoverin (1-3), calcineurin B, troponin C, visinin, neurocalcin, calmodulin, parvalbumin, and the like. For pH, a system based on hisactophilins may be employed. Hisactophilins are myristoylated histidine-rich 20 proteins known to exist in *Dictyostelium*. Their binding to actin and acidic lipids is sharply pH-dependent within the range of cytoplasmic pH variations. In living cells membrane binding seems to override the interaction of hisactophilins with actin filaments. At pH 6.5 they locate to the plasma membrane and nucleus. In contrast, at pH 7.5 they evenly distribute throughout the cytoplasmic space. This change of distribution is reversible and 25 is attributed to histidine clusters exposed in loops on the surface of the molecule. The reversion of intracellular distribution in the range of cytoplasmic pH variations is in accord with a pK of 6.5 of histidine residues. The cellular distribution is independent of myristoylation of the protein. By fusing FPs (Fluorescent Proteins) to hisactophilin the intracellular distribution of the fusion protein can be followed by laser scanning, confocal 30 microscopy or standard fluorescence microscopy. Quantitative fluorescence analysis can be done by performing line scans through cells (laser scanning confocal microscopy) or other electronic data analysis (e.g., using metamorph software (Universal Imaging Corp) and averaging of data collected in a population of cells. Substantial pH-dependent redistribution of hisactophilin-FP from the cytosol to the plasma membrane occurs within 35

1-2 min and reaches a steady state level after 5-10 min. The reverse reaction takes place on a similar time scale. As such, hisactophilin-fluorescent protein fusion protein that acts in an analogous fashion can be used to monitor cytosolic pH changes in real time in live mammalian cells. Such methods have use in high throughput applications, e.g., in the measurement of pH changes as consequence of growth factor receptor activation (e.g. epithelial or platelet-derived growth factor) chemotactic stimulation/ cell locomotion, in the detection of intracellular pH changes as second messenger, in the monitoring of intracellular pH in pH manipulating experiments, and the like. For detection of PKC activity, the reporter system exploits the fact that a molecule called MARCKS (myristoylated alanine-rich C kinase substrate) is a PKC substrate. It is anchored to the plasma membrane via myristoylation and a stretch of positively charged amino acids (ED-domain) that bind to the negatively charged plasma membrane via electrostatic interactions. Upon PKC activation the ED-domain becomes phosphorylated by PKC, thereby becoming negatively charged, and as a consequence of electrostatic repulsion MARCKS translocates from the plasma membrane to the cytoplasm (called the "myristoyl-electrostatic switch"). Fusion of the N-terminus of MARCKS ranging from the myristoylation motif to the ED-domain of MARCKS to fluorescent proteins of the present invention makes the above a detector system for PKC activity. When phosphorylated by PKC, the fusion protein translocates from the plasma membrane to the cytosol. This translocation is followed by standard fluorescence microscopy or confocal microscopy e.g. using the Cellomics technology or other High Content Screening systems (e.g. Universal Imaging Corp./Becton Dickinson). The above reporter system has application in High Content Screening, e.g., screening for PKC inhibitors, and as an indicator for PKC activity in many screening scenarios for potential reagents interfering with this signal transduction pathway. Methods of using fluorescent proteins as biosensors also include those described in U.S. Patent Nos. 972,638; 5,824,485 and 5,650,135 (as well as the references cited therein) the disclosures of which are herein incorporated by reference.

The subject fluorescent proteins also find use in applications involving the automated screening of arrays of cells expressing fluorescent reporting groups by using microscopic imaging and electronic analysis. Screening can be used for drug discovery and in the field of functional genomics: e.g., where the subject proteins are used as markers of whole cells to detect changes in multicellular reorganization and migration, e.g., formation of multicellular tubules (blood vessel formation) by endothelial cells, migration of cells through Fluoroblok Insert System (Becton Dickinson Co.), wound healing, neurite outgrowth, etc.; where the proteins are used as markers fused to peptides (e.g., targeting sequences) and proteins that allow the detection of change of intracellular location as indicator for cellular activity, for example: signal transduction, such as kinase and transcription factor translocation upon stimuli, such as protein kinase C, protein kinase A, transcription factor NFkB, and NFAT; cell cycle proteins, such as cyclin A,

cyclin B1 and cyclinE; protease cleavage with subsequent movement of cleaved substrate, phospholipids, with markers for intracellular structures such as endoplasmic reticulum, Golgi apparatus, mitochondria, peroxisomes, nucleus, nucleoli, plasma membrane, histones, endosomes, lysosomes, microtubules, actin) as tools for High Content Screening: co-localization of other fluorescent fusion proteins with these localization markers as indicators of movements of intracellular fluorescent fusion proteins/peptides or as marker alone; and the like. Examples of applications involving the automated screening of arrays of cells in which the subject fluorescent proteins find use include: U.S. Patent No. 5,989,835; as well as WO/0017624; WO 00/26408; WO 00/17643; and WO 00/03246; the disclosures of which are herein incorporated by reference.

The subject fluorescent proteins also find use in high through-put screening assays. The subject fluorescent proteins are stable proteins with half-lives of more than 24h. Also provided are destabilized versions of the subject fluorescent proteins with shorter half-lives that can be used as transcription reporters for drug discovery. For example, a protein according to the subject invention can be fused with a putative proteolytic signal sequence derived from a protein with shorter half-life, e.g., PEST sequence from the mouse ornithine decarboxylase gene, mouse cyclin B1 destruction box and ubiquitin, etc. For a description of destabilized proteins and vectors that can be employed to produce the same, see e.g., U.S. Patent No. 6,130,313; the disclosure of which is herein incorporated by reference. Promoters in signal transduction pathways can be detected using destabilized versions of the subject fluorescent proteins for drug screening, e.g., AP1, NFAT, NFkB, Smad, STAT, p53, E2F, Rb, myc, CRE, ER, GR and TRE, and the like.

The subject proteins can be used as second messenger detectors, e.g., by fusing the subject proteins to specific domains: e.g., PKCgamma Ca binding domain, PKCgamma DAG binding domain, SH2 domain and SH3 domain, etc.

Secreted forms of the subject proteins can be prepared, e.g. by fusing secreted leading sequences to the subject proteins to construct secreted forms of the subject proteins, which in turn can be used in a variety of different applications.

The subject proteins also find use in fluorescence activated cell sorting applications. In such applications, the subject fluorescent protein is used as a label to mark a population of cells and the resulting labeled population of cells is then sorted with a fluorescent activated cell sorting device, as is known in the art. FACS methods are described in U.S. Patent Nos. 5,968,738 and 5,804,387; the disclosures of which are herein incorporated by reference.

The subject proteins also find use as in vivo marker in animals (e.g., transgenic animals). For example, expression of the subject protein can be driven by tissue specific promoters, where such methods find use in research for gene therapy, e.g., testing

efficiency of transgenic expression, among other applications. A representative application of fluorescent proteins in transgenic animals that illustrates this class of applications of the subject proteins is found in WO 00/02997, the disclosure of which is herein incorporated by reference.

Additional applications of the subject proteins include: as markers following injection into cells or animals and in calibration for quantitative measurements (fluorescence and protein); as markers or reporters in oxygen biosensor devices for monitoring cell viability; as markers or labels for animals, pets, toys, food, etc.; and the like.

The subject fluorescent proteins also find use in protease cleavage assays. For example, cleavage inactivated fluorescence assays can be developed using the subject proteins, where the subject proteins are engineered to include a protease specific cleavage sequence without destroying the fluorescent character of the protein. Upon cleavage of the fluorescent protein by an activated protease fluorescence would sharply decrease due to the destruction of a functional chromophor. Alternatively, cleavage activated fluorescence can be developed using the subject proteins, where the subject proteins are engineered to contain an additional spacer sequence in close proximity/or inside the chromophor. This variant would be significantly decreased in its fluorescent activity, because parts of the functional chromophor would be divided by the spacer. The spacer would be framed by two identical protease specific cleavage sites. Upon cleavage via the activated protease the spacer would be cut out and the two residual "subunits" of the fluorescent protein would be able to reassemble to generate a functional fluorescent protein. Both of the above types of application could be developed in assays for a variety of different types of proteases, e.g., caspases, etc.

The subject proteins can also be used in assays to determine the phospholipid composition in biological membranes. For example, fusion proteins of the subject proteins (or any other kind of covalent or non-covalent modification of the subject proteins) that allows binding to specific phospholipids to localize/visualize patterns of phospholipid distribution in biological membranes also allowing colocalization of membrane proteins in specific phospholipid rafts can be accomplished with the subject proteins. For example, the PH domain of GRP1 has a high affinity to phosphatidyl-inositol tri-phosphate (PIP3) but not to PIP2. As such, a fusion protein between the PH domain of GRP1 and the subject proteins can be constructed to specifically label PIP3 rich areas in biological membranes.

Yet another application of the subject proteins is as a fluorescent timer, in which the switch of one fluorescent color to another (e.g. green to red) concomitant with the ageing of the fluorescent protein is used to determine the activation/deactivation of gene expression, e.g., developmental gene expression, cell cycle dependent gene expression, circadian rhythm specific gene expression, and the like

The antibodies of the subject invention, described above, also find use in a number of applications, including the differentiation of the subject proteins from other fluorescent proteins.

5 KITS

Also provided by the subject invention are kits for use in practicing one or more of the above described applications, where the subject kits typically include elements for making the subject proteins, e.g., a construct comprising a vector that includes a coding region for the subject protein. The subject kit components are typically present in a
10 suitable storage medium, e.g., buffered solution, typically in a suitable container. Also present in the subject kits may be antibodies to the provided protein. In certain embodiments, the kit comprises a plurality of different vectors each encoding the subject protein, where the vectors are designed for expression in different environments and/or under different conditions, e.g., constitutive expression where the vector includes a strong promoter for expression in mammalian cells, a promoterless vector with a multiple cloning
15 site for custom insertion of a promoter and tailored expression, etc.

In addition to the above components, the subject kits will further include instructions for practicing the subject methods. These instructions may be present in the subject kits in a variety of forms, one or more of which may be present in the kit. One form
20 in which these instructions may be present is as printed information on a suitable medium or substrate, e.g., a piece or pieces of paper on which the information is printed, in the packaging of the kit, in a package insert, etc. Yet another means would be a computer readable medium, e.g., diskette, CD, etc., on which the information has been recorded. Yet another means that may be present is a website address which may be used via the
25 internet to access the information at a removed site. Any convenient means may be present in the kits.

The following examples are offered by way of illustration and not by way of limitation.

30 EXPERIMENTAL

I. Wild-Type Anthozoan Proteins

35 The following table summarizes the properties of nine specific wild type Anthozoan proteins of the subject invention:

TABLE I

NFP	Species	Identifier	Absorb. Max. Nm	Emission Max. Nm	Maximum Extinction Coeff.	Relative Quantum Yield*	Relative Brightness**
1	Anemonia majano	amFP486	458	486	40,000	0.3	0.43
2	Clavularia sp.	cFP484	456	484	35,300	0.6	0.77
3	Zoanthus sp.	zFP506	496	506	35,600	0.79	1.02
4	Zoanthus sp.	zFP538	528	538	20,200	0.52	0.38
5	Discosoma striata	dsFP483	443	483	23,900	0.57	0.50
6	Discosoma sp. "red"	drFP583	558	583	22,500	0.29	0.24
7	Anemonia sulcata	asFP600	572	596	56,200	<0.001	-
8	Discosoma sp "green"	dgFP512	502	512	20,360	0.3	0.21
9	Discosoma sp. "magenta"	dmFP592	573	593	21,800	0.11	0.09

*relative quantum yield was determined as compared to the quantum yield of *A. victoria* GFP.

**relative brightness is extinction coefficient multiplied by quantum yield divided by the same value for *A. victoria* GFP.

The sequences of the above wild type proteins and cDNAs encoding the same are provided in Figures 1 to 9.

II. amFP486 Characterization and Mutants Thereof (NFP-1; AmCyan)

A. Construction of amFP486 Mutants

Two mutants of amFP486 were generated, Mut15 and Mut32. Mut32 has the following amino acid substitutions relative to the wildtype: K65M according to GFP alignment numbering as described in Matz et al., supra, or K68M according to self numbering. Mut15 has the following amino acid substitutions relative to the wildtype: K65L according to GFP alignment numbering as described in Matz et al., supra, or K68L according to self numbering. Compared with wildtype amFP486 nucleic acid sequence, Mut15 has the following point mutations: A101G, T129C, AAA 202-204TTG, and C240T. (i.e., A to G at position 101 (numbered from beginning of ATG), T to C at position 129; AAA to TTG at positions 202-204; C to T at position 240).

Table 2 lists the spectral properties of Mut15 and Mut32.

TABLE 2**Spectral Properties of the Isolated Mut15 and Mut32**

Species	NFP Name	Absorbance Maximum nm	Emission Maximum nm	Maximum Extinction Coeff.	Quantum Yield	Relative Brightness *
Anemonia majano	Mut15	460	485	53,400	0.32	0.78
Anemonia majano	Mut32	466	488	36,000	0.42	0.69

*relative brightness is extinction coefficient multiplied by quantum yield divided by the same value for *A. victoria* GFP.

In addition, a non aggregating mutant of Mut 32 was prepared, where this mutant had the following substitutions: K6E, K68M as compared to wild type. The orf nucleotide and amino acid sequences of this mutant are provided in Figure 10.

Additional mutants of NFP-1 include (according to GFP numbering): Y66H; Y66W; A167M; and A167I.

B. Construction and Functional Analysis of Vectors

Mut32 DNA was amplified via PCR and reconstructed to EGFP-N1 backbone with BamHI and NotI restriction enzyme sites. This vector has the same multiple cloning sites as EGFP-N1 (Clontech Laboratories, Palo Alto, CA).

Functional tests of the generated vectors were performed by transient transfection in 293 cells. After 24-hour expression, brighter fluorescent intensity and less photobleaching of pCNFPMut32-N1 were observed by microscopy when compared with pECFP-N1 side by side.

Mut32 has fast folding and bright fluorescent intensity, which makes it useful for number of applications. Some fusion proteins were tested, such as PKC-gamma-CNFP. PKC was observed to translocate from cytosol to the plasma membrane when cells were treated with PMA (phorbol 12-myristate 13-acetate).

C. Generation of Destabilized amFP486 Vectors as Transcription Reporters

Three destabilized amFP486 vectors were constructed by fusing different mouse ODC degradation domains such as d1, d2 and d376 to the C-terminal of wild type amFP486 as described in U.S. Patent No. 6,306,600; the disclosure of which is herein incorporated by reference. The vectors were constructed in the EGFP-N1 backbone.

Vectors of pCRE-d1CNFP and pNF- κ B-d1CNFP were constructed by placing d1CNFP downstream of cAMP response element (CRE) or NF- κ B response element, respectively. Expression of d1CNFP is up-regulated upon activation of these response elements.

D. Functional Analysis of Destabilized amFP486

Functional tests of the destabilized amFP486 were performed by transient transfection in 293 cells. After 24-hour expression, the fluorescent intensity was decreased gradually from d2, d1 and d376 because of the fusion with different mouse ODC degradation domains. After 4-hour treatment with protein synthesis inhibitor cycloheximide, d2 fluorescent intensity did not change very much; however, d1 fluorescent intensity decreased further 50% of its original intensity. The half-life of d1 is around 4 hours.

MODCd1 is a valuable tool for application as a transcription reporter. However, compared with EGFP-d1 (1-hour half-life), pCNFP-MODCd1 half-life (4 hours) is still long, so further mutagenesis for MODC degradation domain is still needed for shorter half-life version.

Functional tests of vectors pCRE-d1CNFP and pNF- κ B-d1CNFP were performed by transient transfection in HEK 293 cells. 16 hours post transfection, 10 μ m forskolin was added to induce CRE and 100 ng/ml TNF-alpha was added to induce NF- κ B for 6 hours. Expression of d1CNFP was analyzed using FACS Calibur. Up to 7 fold increase of fluorescence in forskolin induced CRE activation and 4 fold increase of fluorescence in TNF- α induced NF-KB activation was observed.

E. Construction and Functional Test for Humanized Mut32 (phCNFP-N1)

Since mammalian expression is a very popular tool, a human favored codon version of this mutant is needed for better expression in mammalian cells. To generate humanized Mut32, the Mut 32 sequence was first changed to human favored codon and 23 oligos (12F and 11R) were designed. Next, four rounds of PCR amplification were performed, each round for 20 cycles. PCR cycle was designed as follows: 94°C for 1 min; 94°C for 1 min; 40°C for 1 min; and 72°C for 1 min. The four rounds were: for 1st round, mixing 2 μ l each of every 4 oligos (60 bp), 5 μ l buffer, 1 μ l pfu, 1 μ l dNTP to make total volume of 50 μ l. After 20 cycles of PCR, 5 sets of 150 bp and 1 set of 4 last oligos of 90 bp products were obtained. For 2nd round, mixing new crude PCR products 10 μ l each, 5 μ l buffer, 1 μ l pfu, 1 μ l dNTP to make total volume of 50 μ l. After 20 cycles of PCR, 2 sets of 270 bp and 1 set of 210 bp PCR products were obtained. For 3rd round, mixing new crude PCR products. After 20 cycles of PCR, 1 set of 510 bp and 1 set of 450 bp products were obtained. For 4th round, mixing new crude products. After 20 cycles of PCR, final PCR product (690 bp) was obtained. Further PCR amplification was performed using 1F and 11R primers. As a result, humanized Mut32 was generated. This humanized Mut32 was constituted into EGFP-N1 backbone.

F. Expression of Wildtype and Mutant amFP486 in Mammalian Cells

The original plasmid amFP486 DNAs (wildtype, Mut15 and Mut32 in pQE30) were used to construct N1 version of amFP486 wildtype, Mut15 and Mut32 as described above. The DNAs were inserted into *E.coli* DH5 α . HEK 293 cells were transfected with each of the three N1 constructs using Calcium Phosphate method (Clontech product #K2051-1).

The fluorescent intensity of the transfected cells was analyzed on FACS using FL1 (510/30) detecting channel. Five samples were analyzed in parallel for each construct. The observed mean value of FL1 fluorescent intensity of the M1 population of each sample is summarized in Table 3. It shows that the average of the mean value of each construct (Wildtype, Mut15, and Mut32) has no significant difference.

TABLE 3

FL1 Fluorescent Intensity of M1 Population

Sample #	Wildtype	Mut15	Mut32
1	82.84	106.95	84.51
2	77.52	108.73	91.41
3	111.85	97.08	91.30
4	113.06	90.16	98.16
5	104.95	86.34	111.44
Mean	98.04	97.85	95.36

G. Generation and Expression of Fusion Protein Mut15-mdm2

The Mut15-mdm2 fusion was generated by the following steps: first, mdm2 DNA was obtained by amplifying human Marathon cDNA library (Burke's Lymphoma) using primers:

ATGTGCAATACCAACATGTCTGTACC (SEQ ID No. 19) and CTAGGGGAAATAAGTTAGCAC (SEQ ID No. 20); secondly, the purified PCR product was then amplified with primers:

GGAATTCCAGCCATGGTGTGCAATACCAACATGTCTGTACC (SEQ ID No. 21) and TCCCCCGGGGGGAAATAAGTTAGCAC (SEQ ID No. 22)

in order to add Kozac sequence and restriction sites; thirdly, the purified PCR product from step 2 was digested with EcoR I and Sma I and inserted into EcoR I and SmaI of NFP1Mut15-N1 vector (this vector was generated using BamH I and Not I sites of the pEGFP-N1 backbone). The generated Mut15-mdm2 fusion was then expressed in HEK293 cells.

H. Comparison of the Protein Fluorescent Intensity

pQE30 amFP486 wildtype, Mut15 and Mut32 were transformed into DH5 α . The bacteria grew in the presence of 1 mM IPTG overnight to induce the protein expression.

Cells were lysed in 100 mM Tris, pH8.0 by sonication. Cell lysate was collected after centrifuge at 3000 rpm for 15 minutes at room temperature. The proteins were purified with TALON™ Metal Affinity Resin (Clontech Laboratories, Inc., Palo Alto, CA). Briefly, after the protein was absorbed on the resin, the beads were washed in stepwise with first wash, then first elution (50 mM imidazole) and second elution (200 mM imidazole) in 100 mM Tris-HCl, pH 8.0. The protein is found mostly in the second step elution. It was found that Mut32 has the highest bacterial expression level, while Mut15 has the lowest.

Samples of each elution fraction were run on SDS-PAGE to check the purity of the proteins. Both wildtype amFP486 and Mut32 show a single band, while Mut15 has two more minor bands with higher molecular weight (data not shown).

The protein concentration (fractionII-2) was checked and measured by Bradford assay (Bio-Rad standard assay) using BSA as a standard. The fluorescence intensity (fraction II-2) was determined with a LS50B Luminescence Spectrometer LS50B. EX = 458 nm, EM = 492 nm, both slits = 2.5 nm. Table 4 shows the protein concentration, relative fluorescent (FL) intensity and intensity/μg protein in 700 μl volume. It shows that Mut32 is as bright as wildtype, while Mut15 is worse than the wildtype.

TABLE 4

	Protein Concentration	Relative FL Intensity	Intensity/μg Protein in 700 μl Volume
Wildtype II-2	1.26 μg/5 μl	37.805/5 μl	30.00
Mut15II-2	0.64 μg/5 μl	10.152/5 μl	15.86
Mut32II-2	6.17 μg/5 μl	186.474/5 μl	30.22

III. Characterization of cFP484 and mutants thereof (NFP-2)

A. Mutant Generation

Two deletion mutants were generated by two separate PCR reactions: Δ19 cFP484 lacks the N-terminal first 19 amino acids of cFP484, and Δ38 cFP484 lacks the N-terminal first 38 amino acids of cFP484. Mammalian expression vectors containing the DNA encoding the fluorescent protein Δ19 cFP484 or Δ38 cFP484 are generated, which are named as pΔ19 NFP2-N1 and pΔ38 NFP2-N1, respectively.

B. Transient Expression of Deletion Mutants of cFP484 in Mammalian Cells

HeLa cells were transiently transfected with mammalian expression vector pΔ19 NFP2-N1 which contains the DNA encoding the fluorescent protein Δ19 cFP484. After transfection, cells were incubated for 48 hours at 37°C then fixed in 3.7% formaldehyde. Cells were mounted in mounting medium and observed by fluorescence microscopy. Digital images were taken with MetaMorph software (Universal Imaging Corp.) using a monochrome cooled CCD camera (Roper Scientific). The filter set XF 114 (Omega

Optical) was used to visualize fluorescence emitted by $\Delta 19$ cFP484. The image was pseudocolored. $\Delta 38$ cFP484 is also fluorescent when expressed in HeLa cells.

IV. Characterization of zFP506 and Mutants thereof (NFP-3; ZsGreen)

A. Mutant generation

A mutant of zFP506 was generated, N66M (N65M if numbered according to GFP homology alignment). Compared with wild type zFP506, N66M has the mutation of from "AAC" to "ATG" which results in the corresponding amino acid change from Asparagine (N) to Methionine (M) at the position of 66. The spectral properties of N66M are listed in Table 5.

TABLE 5

Spectral Properties of the Isolated N66M

Species	NFP Name	Absorbance Maximum Nm	Emission Maximum nm	Maximum Extinction Coeff.	Quantum Yield	Relative Brightness *
Zoanthus sp.	N65M	496	506	62,000	0.63	1.78

*relative brightness is extinction coefficient multiplied by quantum yield divided by the same value for *A. victoria* GFP.

The following additional mutants were also generated: FP3-NA: K5E, K10E, N66M (non-aggregating mutant)(See Figure 11); Yellow (A64G, N66K, N69D)(See Figure 12); Yellow/bright (A64G, N66K, N69D, D94N, M120V, K157R, P231S) (See Figure 13).

Additional mutants of NFP-3 include (according to GFP numbering): M167A; M167H; and N65K/N68D.

B. Construction and Functional Analysis of Vectors

Non-humanized zFP506 DNA was amplified via PCR and reconstructed into EGFP-N1 backbone. This vector has the same multiple cloning sites as EGFP-N1. Functional test of the generated vector was performed by transient transfection in 293 cells. 24 hours post transfection, expression of zFP506 was examined under fluorescent microscope. zFP506 showed good fluorescent intensity and comparable to EGFP-N1.

C. Generation of Destabilized zFP506 Vectors as Transcription Reporters

Since zFP506 is very stable, it is necessary to generate destabilized versions of zFP506 in order to observe the rapid turnover of the protein. By using the same technology for destabilized EGFP, two destabilized zFP506 vectors were constructed by fusing mouse ODC degradation domain to the C-terminal of zFP506. The d1 version of destabilized zFP506 has three E to A mutations within MODC degradation domain

comparing to d2 version, therefore result in a shorter half-life of the protein to which MODC degradation domain fused to. Destabilized d1zFP506 and d2zFP506 were constructed in EGFP-N1 backbone

5 D. Functional Analysis of Destabilized zFP506

Wildtype d1zFP506 was transiently transfected into 293 cells. 24 hours after transfection, CHX was added to stop protein synthesis. After 4 hour treatment, cells were examined under fluorescent microscope. It shows that fusion of MODC domain to the zFP506 slightly decreases the fluorescent intensity compared to zFP506 itself. After 4
10 hour treatment, there is 50% fluorescent intensity decrease.

E. Application of Destabilized d1zFP506 as Transcription Reporters

Destabilized d1zFP506 was constructed into pCRE-d1GNFP and pNF- κ B-d1GNFP vectors. Its expression was regulated under cAMP response element (CRE) or NF- κ B response element, respectively. These vectors were transiently transfected into 293 cells, and 24 hours post transfection, the expression of d1GNFP was induced by Forskolin or TNF- α . 6 hours after induction, the culture was analysed by FACS. CRE-d1GNFP showed 7 fold of induction in fluorescence intensity, while 4 fold of induction was obtained in NF- κ B-d1GNFP (data not shown). This demonstrated that the destabilized form of
20 GNFP is applicable as transcription reporters.

F. Construction and Functional Test for Humanized zFP506 and Humanized N66M

Since mammalian expression is a very popular tool, human favored codon version is needed for better expression in mammalian cells. Each piece of human favored codon
25 oligos was linked to form the full length of wild type and/or mutant zFP506 (hGNFP-zFP506; hGNFP-N65M. This humanized zFP506 was constituted into EGFP-N1 backbone.

V. Characterization of zFP538 and mutants thereof (NFP-4; ZsYellow)

30 A mutant of zFP538 M129V (as measured from the start of the protein) was generated. M129V (M128V using GFP numbering) was generated by introducing a wrong nucleotide in PCR during site-specific mutagenesis at position 65. One bright yellow colony was obtained, and the sequence of this clone was performed. It showed that this
35 clone contained wild type amino acid Lysine (K) at position 65, but had a substitution from Methionine (M) to Valine (V) at position 129 (numbering from start of protein; at position 128 if numbering according to GFP homology alignment).

Further investigations showed that M129V has spectral characteristics very similar to wild type protein zFP538 but folds much faster. Table 6 lists the spectral properties of M129V.

TABLE 6

Spectral Properties of the Isolated M128V

Species	NFP Name	Absorbance Maximum nm	Emission Maximum nm	Maximum Extinction Coeff.	Quantum Yield	Relative Brightness *
Zoanthus sp.	M128V	531	540	25,360	0.43	0.50

*relative brightness is extinction coefficient multiplied by quantum yield divided by the same value for *A. victoria* GFP.

The following additional mutants were also generated: FP4-NA (K5E, K9T, M129V)(Non-Aggregating Mutant); Green (K65M GFP numbering; K66M self numbering).

Additional mutants of NFP-1 include (according to GFP numbering): D68N.

A. Construction and Functional Analysis of Vectors

Both wildtype (wt) and mutant zFP538 DNA were amplified via PCR and reconstructed to EGFP-N1 backbone. This vector has the same multiple cloning sites as EGFP-N1. Both pYNFPwt and pYNFPW129V keep the same multiple cloning sites as EGFP-N1. Functional test of the generated vectors was performed by transient transfection in 293 cells. After 24-hour expression, pYNFPwt, pYNFPM129V and EYFP were compared side by side: pYNFPwt showed less fluorescent intensity than EYFP (data not shown); however, pYNFPM129V showed as bright fluorescent intensity as EYFP by fluorescent microscopy.

B. Generation of Destabilized zFP538 Vectors as Transcription Reporters

By using the same technology for destabilized EGFP, destabilized zFP538 vectors were constructed by fusing different mouse ODC degradation domains such as d1 and d2 to the C-terminal of zFP538. The d1 version of destabilized YNFP has three E to A mutations within MODC degradation domain compared to d2 version. Vectors pYNFPM128V-MODCd1 and pYNFPM128V-MODCd2 were constructed in EGFP-N1 backbone.

C. Functional Analysis of Destabilized zFP538

Functional test of the destabilized zFP538 was performed by transient transfection in 293 cells. After 24-hour expression, the fluorescent intensity was decreased gradually from d2 and d1 because of the fusion with different mouse ODC degradation domains.

After 4-hour treatment with protein synthesis inhibitor cycloheximide, d2 fluorescent intensity did not change very much; however, d1 fluorescent intensity decreased further 50% of its original intensity. The half-life of d1 is around 4 hours.

M129V has fast folding and bright fluorescent intensity, which makes it useful for number of applications. Some fusion proteins were tested such as PKC-gamma-YNFP (M129V). PKC-gamma was observed to translocate from cytosol to the plasma membrane when cells were treated with PMA (Phorbol 12-Myristate 13-Acetate).

D. Construction and Functional Test for Humanized M129V

Humanized M129V was generated, and then placed into the pEGFP-N1 backbone. This vector has the same multiple cloning sites as pEGFP-N1. Construction of C1 and pEGFP is in the process.

E. Structural Characterization of Green Mutant

1. analytical ultracentrifugation:

Data show zFP538 is non-dissociating trimer of 80kD/dimer-tetramer equilibrium with S values of about 6 and 8; results not conclusive, but sure that there is more than one species.

2. Structure:

Tetramer, extremely similar to DsRed; no idea why wildtype is yellow, chromophore is identical to that of GFP; although the tetramers of dsRed and zFP538 are essentially identical in the overall configuration and in the positioning of the interfacial regions, the residues involved in interfacial contacts are not conserved. This finding indicates that a lot of variability is tolerated and monomers can be created.

VI. Characterization of dsFP483 and mutants thereof (NFP-5)

Mutants of NFP-5 include (according to GFP numbering): N68S; I112S; and N68S/I112S.

VII. Characterization of drFP583 and mutants thereof (NFP-6; DsRed; DsRed2)

A. Expression in Mammalian Cells

HeLa cells were transfected either with plasmid pDsRed1-N1 (vector containing the DNA encoding drFP583) or plasmid pEGFP-C1 (encoding EGFP from *Aequorea victoria*). Immediately after the transfection, cells were mixed and plated on coverslips. Cells were

incubated for 48 hours at 37°C then fixed in 3.7 % formaldehyde. Cells were mounted in mounting medium and observed by fluorescence microscopy. Images were taken from the same field of view with Chroma filter set 31001 for EGFP and filter set 31002 for drFP583 using a cooled CCD camera (Roper Scientific) and MetaMorph software (Universal Imaging). The images were pseudocolored and overlaid. Phase contrast was taken from the same field of view and overlaid.

B. Generation of Humanized drFP583

Since mammalian expression is a very popular tool, human favored codon version is needed for better expression in mammalian cells. Humanized drFP583 was therefore generated by changing wild type drFP583 nucleotide sequence to optimize the codons for expression of the fluorescent protein. The nucleotide sequence of this humanized mutant is provided in Figure 16.

C. Expression of Humanized drFP583 in Mammalian Cells

HeLa cells were transiently co-transfected with plasmids pECFP-Nuc, pEYFP-Tub and pDsRed1-Mito (humanized drFP583). After transfection, cells were incubated for 48 hours at 37 °C then fixed in 3.7 % formaldehyde. Cells were mounted in mounting medium and observed by fluorescence microscopy. Images were taken of one cell co-expressing all three fluorescent proteins with Omega filter set XF 35 for DsRed1-Mito, XF 104 for EYFP-Tub and XF 114 for ECFP-Nuc using a cooled CCD camera (Roper Scientific) and MetaMorph software (Universal Imaging). Individual images were pseudocolored and overlaid to show all three signals in one image. Protein DsRed1-Mito localizes to mitochondria, EYFP-Tub localizes to the microtubular network, and ECFP-Nuc localizes to the nucleus.

As a conclusion, drFP583 does emit to a low extent also in the cyan (ECFP), green (EGFP) and yellow-green (EYFP) emission channels (filter sets). High expression levels or highly concentrated protein in intracellular structures can therefore result in high signal intensities that will give some bleedthrough in the other emission wavelengths. The bleedthrough is small and should not affect multiple labeling recording in most cases.

D. Mutants of Humanized drFP583

Mutants of humanized drFP583 were generated using error prone PCR technique (Clontech). Mutations occurred at amino acids 42, 71, 105, 120, 161 and 197 (numbering starting from the first Methionine). Table 7 lists the mutants that were generated and their properties.

TABLE 7**Mutants of Humanized drFP583**

Mutant	Mutations	Properties
E5	V105A, S197T	Overnight in <i>E. coli</i> emitting green fluorescence; <i>in vitro</i> maturing to red over 28h at 37°C on 80% (retains 20% green peak); folding faster than wild type drFP583 (~28h)
E8	N42H	Always two peaks green & red (~1:1) folding faster than E5 (~8h)
E5up	V105A	red from the beginning; folding faster than E5 (~12h)
E5 down	S197T	phenotype is similar to E5
E57	V105A, I161T, S197A	like E5 but folding faster (~8-10h); ~5% of green peak left at the end (See Figure 18)
AG4	V71M, V105A, S197T	bright green, no red at all; fast folding (~ 16h)
AG45	V71M, V105A, Y120H, S197T	like AG4 but twice brighter
FP6 (E57)-NA:	R2A, K5E, K9T, V105A, I161T, S197A.	Non-aggregating (See Figure 19)
E5-NA:	R2A, K5E, K9T, V105A, S197T	(non-aggregating Fluorescent Timer) (See Figure 17)
E83	N42H, V71A, I180V	

E. Characterization and Applications of E5 Mutant

E5 (V105A, S197T) changes its fluorescence from green to red over time both *in vitro* and *in vivo*, in *E. coli* and in mammalian cells. Also, E5 folds faster than wild type drFP583 both in *E. coli* and mammalian cells.

Since it allows the “two color” reporting mode for monitoring of the promoter activity, i.e., for both active or shutdown state of the promoter, there is a separate color, serving as an indicator of that state, E5 can be used as a transcriptional reporter. Different from “one color” mode, “two color” mode has a measurable signal (color) present for both states of the process as opposed to “one color” mode (e.g. destabilized GFP) wherein the absence of the color is an indicator of the second state. Namely, newly produced E5 protein fluoresces in green, indicating on-going promoter activity. Over time, the protein will mature, acquiring the red fluorescence. So if the promoter is no longer active, all the protein will eventually mature, resulting in the dominant red fluorescence. In case the promoter is still active both red and green fluorescence will be readily detected. Thus E5 as a “two color” reporter allows study of gene expression similar to destabilized GFP, but with permanent “signature” of past gene activity in the cells, tissues or entire organism. For example, at the tissue level, E5 may help to distinguish the stem cells from differentiated cells. Providing the promoter is only active in the stem cells, the E5 reporter will label the stem cell population in green and red, the progenitor cells would be labeled

predominantly in red, the terminally differentiated will not fluoresce (due to the titration out of protein during cell division).

E5 can be used for spatial and temporal visualization of newly synthesized vs. accumulated fusion proteins. That is, E5 could function like a fusion tag. Possible applications envisaged at different organizational levels. At the cellular level, E5 may help to visualize and distinguish the newly synthesized proteins in various compartments such as outer membrane, microvillae, ER, Golgi, mitochondria, nuclei, various components of cellular matrix and focal adhesion complexes. At the tissue level, E5 may be helpful in visualizing newly formed vs. preexisting structures e.g. membrane junctions, components of extracellular matrix.

One of the most fascinating applications of E5 seems to be in the study of the mother-daughter relationship during the cell division and migration. A wide horizon is opening in the fields of development as well as in the adult organisms to study the cell migration and differentiation. Allowing visualization of the expression "history", E5 can help to distinguishing between the mother cells where the promoter is actually active vs. the daughter cells containing the accumulated protein but not producing fresh protein anymore. It would enable the study of the cell fates during development and organ remodeling, thus distinguishing between cell migration and cell expansion or differentiation.

In conclusion, E5 is basically applicable to any situation where GFP was previously used. Main advantage is that E5 can track down "the history" of promoter activity or protein localization in cells or tissues. With a better protein stability than GFP, E5 allows longer analysis window (wild type drFP583 is stable for at least 4 weeks in *Xenopus*, while EGFP starts to faint after two weeks).

F. Characterization and Applications of E8 Mutant

E8 (N42H) has two fluorescence maximums, green and red, at all times and it folds much faster than drFP583 (Table 7).

Since it detects both green and red fluorescence simultaneously, E8 may be useful for studying processes related to blood circulation and proteins/cells trafficking. Blood absorbs the green fluorescence; thus only the red fluorescence will be visible while the protein is trafficking in the blood. Both green and red fluorescence could be detected outside the bloodstream making the whole process easy to visualize and record. Monitoring both red and green fluorescence simultaneously may also help to reduce the background fluorescence problems for some tissues or cells.

G. Generation of drFP583/dmFP592 Hybrid Using Shuffling Procedure

Non-humanized wild type coding region fragments from drFP583 and dmFP592 were amplified by PCR (22 cycles, 95°C, 15 sec., 68°C 1 min 20 sec.) using 1 ng of

corresponding bacterial expression plasmids (pQE-30 derivatives with drFP583 and dmFP592 inserts, respectively) as templates. Oligonucleotides

A (ACATGG ATCCAGGTCTTCCAAGAATGTTATC, SEQ ID No. 23),

B (TAGTACTCG AGCCAAGTTCAGCCTTA, SEQ ID No. 24),

5 C (ACATGGATCCAG TTGTTC CAAGAATGTGAT, SEQ ID No. 25), and

D (TAGTACTCGAGGCCATTA CCGCTAATC, SEQ ID No. 26)

were used as primers for amplifying these fragments in a concentration of 0.2 mM.

The PCR products were then purified by QIAquick PCR Purification Kit (QIAGEN).

Afterwards, the purified fragments drFP583 and dmFP592 (300-500 ng each) were

10 digested with restriction endonucleases EcoRI, HindIII and DraI (10 U each)

simultaneously. Reactions were performed in BamHI restriction buffer (NEB)

supplemented BSA for 3h at 37°C. Total reaction volume was 30 ml. Upon completion,

the resulted restriction fragments from each restriction reaction were separated by

electrophoresis in agarose gels (1.5%), cut from gel and purified by QIAquick Gel

Purification Kit (QIAGEN). The resulting sets of the purified restriction fragments from

both drFP583 and dmFP592 were combined together and 50 ng of them were put into

ligation mix (1X T4 DNA ligation buffer, 400 NEB U of T4 DNA ligase) in total volume of

30 ml. The ligation was performed for 3h at room temperature and stopped by heating at 70°C within 20 min.

20 The ligation mixture was then diluted by water ten-folds, and 1 ml of the dilution was preformed for PCR reaction (20 cycles, 95°C, 15 sec. 68°C 1min 20 sec) as template.

Four oligonucleotides A, B, C, and D (SEQ ID Nos. 58-61, respectively) were used

simultaneously as primers for amplifying these fragments in a concentration of 0.1 mM each. After electrophoresis in an agarose gel (1.5%), the target fragment was purified by

25 QIAquick Gel Purification Kit (QIAGEN) and digested with restriction endonucleases

BamHI and XhoI (30-50 U each) simultaneously in BamH I restriction buffer (NEB)

supplemented with BSA for 3h at 37°C. After purification, the resulting fragment was

cloned in pQE-30 plasmid linearized by BamHI and Sall. Ligation reaction was performed

in 1X T4 DNA ligation buffer and 400 NEB U of T4 DNA ligase with a total volume of 20 ml

30 for overnight at 16°C. After transformation of *E.coli* cells by 1/5 of the ligation volume and

incubation on LB/1% agar plates which were supplemented by 100 mg/ml Ampacilin and

0.1 mM IPTG at 37°C for overnight, the resulting *E.coli* colonies were screened visually

under fluorescent microscope using rhodamine filter set. The brightest red colonies were

picked up and placed in 200 ml LB medium with 100 mg/ml of Ampacilin. At OD₆₀₀=0.6,

35 the *E.coli* culture was induced by IPTG (final concentration was 1 mM) and the

fermentation continued for overnight. Purification of recombinant protein containing N-

terminus 6xhis tag was performed using TALON metal-affinity resin according to

manufacturer's protocol.

H. Spectral Properties of drFP583/dmFP592 Hybrid

The emission and excitation spectra for drFP583/dmFP592 hybrid protein are basically the same as for dmFP592. Table 8 lists the spectral properties of drFP583/dmFP592 hybrid protein.

TABLE 8

Spectral Properties of drFP583/dmFP592 Hybrid

nFP Name	Absorbance Maximum nm	Emission Maximum nm	Maximum Extinction Coeff.	Relative Quantum Yield*	Relative Brightness**
drFP583/dmFP592	573	592	35,000	0.24	0.3

*relative quantum yield was determined as compared to the quantum yield of *A. victoria* GFP.

**relative brightness is extinction coefficient multiplied by quantum yield divided by the same value for *A. victoria* GFP.

I. Humanized drFP583/dmFP592 Hybrid and Mutants

drFP583/dmFP592 hybrid was humanized. Further, two mutants were generated based on the humanized drFP583/dmFP592, i.e., drFP583/dmFP592-2G and drFP583/dmFP592-Q3. drFP583/dmFP592-2G (i.e. 6/92G) contains two substitutions, K15Q and T217S (Self numbering). This mutant was derived from the humanized drFP583/dmFP592 hybrid gene by random mutagenesis using Diversity PCR Mutagenesis Kit (Clontech) according to the corresponding protocol. drFP583/dmFP592-Q3 (6/9Q) contains three substitutions, K15Q and K83M and T217S (self numbering). drFP583/dmFP592-Q3 mutant was derived from drFP583/dmFP592-2G mutant by random mutagenesis using Diversity PCR Mutagenesis Kit (Clontech) according to the corresponding protocol. A non-aggregating mutant of 6/9 Q, i.e. 6/9 QNA, was also produced, which the following substitutions: K5E, K9T, K15Q, K83M, T217S (self numbering). The sequences of a humanized 6/9 Q hybrid protein are provided in Figure 22.

drFP583/dmFP592-2G has similar brightness and folding rate as for non-humanized drFP583/dmFP592 hybrid. While drFP583/dmFP592-Q3 could be seen in *E. coli* cells as more dark red than parental variant, i.e., drFP583/dmFP592-2G, and the purified protein solution has purple color. drFP583/dmFP592-Q3 has the emission maximum of 616 nm and excitation maximum of 580 nm.

J. Applications of Hybrid Mutants

Similar to fluorescent protein drFP583 or dmFP592, drFP583/dmFP592-Q3 can be used as a tool for investigation of protein expression, transport and protein interactions *in vivo*, monitoring of promoter activity, and as a transcription reporter or fusion tag.

Besides, drFP583/dmFP592-Q3 can be chosen as the most convenient partner to one of the existing green fluorescent protein variants in two/triple color labeling assays for simultaneous detection of expression of two or more proteins *in vivo* due to its strongly red-shifted position of emission maximum and practical absence of excitation in green part of spectrum except any spectral overlapping and background fluorescence.

The method of generating drFP583/dmFP592 hybrid can have a general utility for generating hybrid genes (i.e., genes containing parts of different genes in various combinations) with improved fluorescent characteristics.

Additionally, drFP583/dmFP592-Q3 is the first red-shifted mutant, which demonstrates that spectral-shifted mutant could be obtained by random mutagenesis.

K. Further Characterization.

Mutant name	Position of mutation(s)	Parental clone/ gene	residual green fluorescence (%) from red)		in vitro maturation t _{1/2}
DsRed	wild type		4-5 after 18-24 hours	Not seen ON in Ecoli, 16-24 h in 293 cells	9-10
E5	V105A,S197T,	FP#6 (w.t.)	20% after 43 hours, eventually drops to 10%	In <i>E.coli</i> GREEN o/night, in 293 cells both color develop faster than wt	9-10
E5-UP	V105A	E5(split)	3% after 24 hours eventually zero	better folding in bacteria	9-10
E5-DOWN	S197T	E5(split)	similar to E5	similar to E5	9-10
E57	V105A,I161T, S197A	E5	5% after 8h, 4% after 24 hours	better folding and faster maturation of fluorophore	3-4
AG4	V71M, V105A,S197T	E5	100% green at all time	bright GREEN, no RED at all	1.5-2
AG45	V71M,V105A, Y120H,S197T	AG4	100% green at all time	~2x brighter than AG4	1.5-2
E8	N42H	FP#6 (w.t.)	Green/Red ratio is 3:2	better folding and faster maturation of fluorophore	3-4
E83	N42H,V71A, I180V	E8	Green/Red ratio is 1:1	same as E8	3-4

green fluorescence:

RFP about 4-5% wt after 18-24 hours, remains.

E5 about 20% after 43 hours eventually drops to 10%

E5up about 3% after 24hours eventually goes to zero.

E5down about 20% after 24 hours, eventually drops to 10%

E57 5% after 8 hours, 4% after 24 hours; faster maturation than wt RFP

VIII. Characterization of asFP600 and Mutants thereof (NFP-7; AsRed)

A. Mutant Generation

A mutant of asFP600 was generated, Mut1. Compared with wild type asFP600, Mut1 has the following substitutions: T68A and A143S from start of protein (T to A at position 70 and A to S at position 148 under numbering according to GFP). Target substitution A143S was generated by means of site-specific mutagenesis using PCR with primers that carried the mutation. During this mutagenesis random substitution T68A was generated by introducing a wrong nucleotide in PCR. The substitution T68A is not necessary for fluorescence and practically does not affect the fluorescence. Table 9 lists the spectral properties of Mut1. Another mutant of asFP600 was generated, having a substitution of the Alanine at position 184 to Serine (according to GFP numbering).

TABLE 9

Spectral Properties of the Isolated Mut1

Species	NFP Name	Absorbance Maximum Nm	Emission Maximum nm	Maximum Extinction Coeff.	Quantum Yield	Relative Brightness *
Anemonia sulcata	Mut1	573	595	15,500	0.05	0.03

*relative brightness is extinction coefficient multiplied by quantum yield divided by the same value for *A. victoria* GFP.

Yet one more mutant was generated, i.e., mut M35-5 /mut1(=7A): F4L, K12R, F35L, T68A, F84L, A143S, K163E, M202L (positions relative to the start of the protein)

A non-aggregating version of this mutant was also made, i.e., 7A-NA: K6T, K7E, F4L, K12R, F35L, T68A, F84L, A143S, K163E, M202L (positions relative to the start of the protein)

The humanized FP-7 sequences are provided in Figure 21.

B. Construction and Functional Analysis of Vectors

Non-humanized mutant asFP600 (RNFP) DNA were amplified via PCR and reconstructed to EGFP-N1 backbone. This vector (pRNFP-N1) has the same multiple cloning sites as EGFP-N1.

Functional test of the generated vector was performed by transient transfection in 293 cells. 24 hours post transfection, expression of asFP600 was examined under

fluorescent microscope. asFP600 showed good fluorescent intensity, however, the expression of asFP600 concentrated at the nucleus.

C. Generation of Cytosal Expressed asFP600

Since the nuclear localization of asFP600 limited some of the application of this protein as transcription reporter or pH sensor, cytosol expression of this protein would be needed for this purpose. A nuclear export sequence in humanized codon usage was fused to the N-terminus of asFP600, and placed into the EGFP-N1 vector, resulted in pNE-RNFP.

Functional test of NE-RNFP is performed by transiently transfect the pNE-RNFP into 293 cells. 24 hours post transfection, expression of NE-RNFP is examined under fluorescence microscope. Red fluorescence was observed to be distributed in the cytosol but not in the nucleus.

D. Generation of Destabilized asFP600 Vectors as Transcription Reporters

Since asFP600 is very stable, it is necessary to generate destabilized versions of asFP600 in order to observe the rapid turnover of the protein. By using the same technology for destabilized EGFP, two destabilized NE-RNFP vectors were constructed by fusing mouse ODC degradation domain to the C-terminal of NE-RNFP. The d1 version of destabilized RNFP has three E to A mutations within MODC degradation domain comparing to d2 version, therefore result in a shorter half-life of the protein to which MODC degradation domain fused. Destabilized d1RNFP and d2RNFP were constructed in EGFP-N1 backbone.

E. Functional Analysis of Destabilized asFP600

d2 version of the none-humanized asFP600 was transiently transfected into 293 cells. One day after transfection, CHX was added to inhibit protein synthesis. 3 hours after treatment, cells were examined under fluorescent microscope. It showed that fluorescent intensity decreased ~50%.

F. Construction and Functional Test for Humanized Mut1

Humanized Mut1 was generated. The humanized Mut1 was then placed into the pEGFP-N1 backbone. This vector has the same multiple cloning sites as pEGFP-N1. Construction of C1 and pEGFP is in the process.

It is evident from the above discussion and results that the subject invention provides important new chromoproteins and fluorescent proteins and nucleic acids encoding the same, where the subject proteins and nucleic acids find use in a variety of

different applications. As such, the subject invention represents a significant contribution to the art.

IX. Summary Table of Mutant Fluorescent Proteins

Table 10

NFP#	name	Mutants: aa numbering according to GFP homology alignment	Mutants: aa numbering according to aa sequence of protein proper
1	amFP486	K65M (mut32) K65L (mut15) FP1-NA: K6E, K65M (non-aggregating mut 32)	K68M (mut32) K68L (mut15) FP1-NA: K6E, K68M (non-aggregating mut 32)
2	CFP484	19 (N-terminal 19 aa deleted) 38 (N-terminal 38 aa deleted)	
3	zFP506	Mut: N65M FP3-NA: K5E, K10E, N65M (non- aggregating mutant) Yellow (A63G, N65K, N68D) Yellow/bright (A63G, N65K, N68D, D95N, M119V, K158R, P232S)	Mut: N66M FP3-NA: K5E, K10E, N66M (non- aggregating mutant) Yellow (A64G, N66K, N69D) Yellow/bright (A64G, N66K, N69D, D94N, M120V, K157R, P231S)
4	zFP540	Mut: M128V FP4-NA: K5E, K9T, M128V (non- aggregating mutant) Green (K65M)	Mut: M129V FP4-NA: K5E, K9T, M129V (non- aggregating mutant) Green (K66M)
5	dsFP483		
6	DsFP583		FP6 (E57)-NA: R2A, K5E, K9T, V105A, I161T, S197A. E5 (V105A, S197T; Fluorescent Timer) E5-NA: NA: R2A, K5E, K9T, V105A, S197T (non-aggregating Fluorescent Timer); E5up (V105A; faster maturing, brighter) E5down (S197T, fluorescent timer phenotype) E57 (V105A, I161T, S197A; faster chromophore maturation, brighter) AG4 (V71M, V105A, S197T; green only) AG45 (V71M, V105A, Y120H, S197T; green only) E8 (N42H; green/red) E83 (N42H, V71A, I180V; green/red)
6/9 hybrid			6/9 Q (K15Q, K83M, T217S) 6/9Q-NA: (S2del, C3del, accidentally deleted) K5E, K9T, K15Q, K83M, T217S. Other mutants: 6/92G (K15Q, T217S)
7 7A	asFP595	mut1: T70A, A148S mut M35-5 /mut1(=7A): F7L, K15R, F38L, T70A, F88L, A148S, K170E, M208L.	mut1: T68A, A143S mut M35-5 /mut1(=7A): F4L, K12R, F35L, T68A, F84L, A143S, K163E, M202L

NFP#	name	Mutants: aa numbering according to GFP homology alignment	Mutants: aa numbering according to aa sequence of protein proper
			7A-NA: : K6T, K7E, F4L, K12R, F35L, T68A, F84L, A143S, K163E, M202L (non-aggregating version of 7A)

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. The citation of any publication is for its disclosure prior to the filing date and should not be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it is readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.